



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 213/74, A61K 31/495, C07D 401/04, 403/04, 243/08, 401/14, 405/04, 409/04, 409/14, 417/04, 417/14, 215/38</p>	A1	<p>(11) International Publication Number: WO 99/21834</p> <p>(43) International Publication Date: 6 May 1999 (06.05.99)</p>									
<p>(21) International Application Number: PCT/DK98/00465</p> <p>(22) International Filing Date: 27 October 1998 (27.10.98)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>1225/97</td> <td>27 October 1997 (27.10.97)</td> <td>DK</td> </tr> <tr> <td>0409/98</td> <td>24 March 1998 (24.03.98)</td> <td>DK</td> </tr> <tr> <td>PA 1998 00796</td> <td>19 June 1998 (19.06.98)</td> <td>DK</td> </tr> </table> <p>(71) Applicant (for all designated States except US): NEUROSEARCH A/S [DK/DK]; Smedeland 26B, DK-2600 Glostrup (DK).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): PETERS, Dan [SE/SE]; Nygatan 17, S-232 31 Ärlöv (SE). OLSEN, Gunnar, M. [DK/DK]; (DK). NIELSEN, Simon, Feldbæk [DK/DK]; (DK). NIELSEN, Elsebet, Østergaard [DK/DK]; NeuroSearch a/s, Smedeland 26B, DK-2600 Glostrup (DK).</p> <p>(74) Common Representative: NEUROSEARCH A/S; Smedeland 26B, DK-2600 Glostrup (DK).</p>		1225/97	27 October 1997 (27.10.97)	DK	0409/98	24 March 1998 (24.03.98)	DK	PA 1998 00796	19 June 1998 (19.06.98)	DK	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
1225/97	27 October 1997 (27.10.97)	DK									
0409/98	24 March 1998 (24.03.98)	DK									
PA 1998 00796	19 June 1998 (19.06.98)	DK									
<p>(54) Title: HETEROARYL DIAZACYCLOALKANES AS CHOLINERGIC LIGANDS AT NICOTINIC ACETYLCHOLINE RECEPTORS</p> <div style="text-align: center; margin: 20px 0;"> <p>(I)</p> </div> <p>(57) Abstract</p> <p>The present invention discloses compounds of formula (I) any of its enantiomers or any mixture thereof, isotopes thereof or a pharmaceutically acceptable salt thereof; wherein n is 1, 2 or 3; m is 0, 1 or 2; R represents hydrogen, alkyl, cycloalkyl, cycloalkylalkyl or aralkyl; and R¹ represents aminophenyl; nitrophenyl; hydroxyphenyl, alkoxyphenyl; a monocyclic 5 to 6 membered heterocyclic group which may be substituted one or more times with substituents or R¹ represents a bicyclic heterocyclic group, composed of 5 to 6 membered monocyclic heterocyclic group fused to a benzene ring, and which may be substituted one or more times with substituents. The compounds of the invention are useful as nicotinic ACh receptor ligands.</p>											

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HETEROARYL DIAZACYCLOALKANES AS CHOLINERGIC LIGANDS AT NICOTINIC ACETYLCHOLINE RECEPTORS

The present invention relates to novel heteroaryl diazacycloalkane derivatives which are cholinergic ligands at nicotinic ACh receptors. The compounds of the invention are useful for
5 the treatment of conditions or disorders or diseases involving the cholinergic system of the central or peripheral nervous system, pain, inflammatory diseases, diseases caused by smooth muscle contractions and as assistance in the cessation of chemical substance abuse.

Background

10

The endogenous cholinergic neurotransmitter, acetylcholine, exert its biological effect via two types of cholinergic receptors; the muscarinic ACh receptors and the nicotinic ACh receptors. As it is well established that muscarinic ACh receptors dominate quantitatively over nicotinic ACh receptors in the brain area important to memory and cognition, much research aimed at
15 the development of agents for the treatment of memory related disorders have focused on the synthesis of muscarinic ACh receptor modulators. Recently, however, an interest in the development of nicotinic ACh receptor modulators has emerged. Several diseases are associated with degeneration of the cholinergic system i.e. senile dementia of the Alzheimer type, vascular dementia and cognitive impairment due to the organic brain damage disease
20 related directly to alcoholism. Indeed several CNS disorders can be attributed to a cholinergic deficiency, a dopaminergic deficiency, an adrenergic deficiency or a serotonergic deficiency. Alzheimer's disease is characterised by a profound loss of memory and cognitive functions caused by a severe depletion of cholinergic neurons, i.e. neurons that release acetylcholine. A reduction in the number of nicotinic ACh receptors are also observed with the progression of
25 Alzheimer's disease. It is believed that the neurons in the cortex that die with the progression of Alzheimer's disease do so because of lack of stimulation of the nicotinic ACh receptors. It is predicted that treatment of Alzheimer's patients with nicotinic ACh receptor modulators will not only improve the memory of patients but in addition act to keep these neurons alive. Smoking actually seems to protect individuals against neurodegeneration and compounds behaving on
30 these receptor may very likely have a generally neuroprotective effect.

However degeneration of the cholinergic system is not limited to individuals suffering from i.e. Alzheimers disease but is also seen in healthy aged adults and rats. Therefore it is suggested that the cholinergic system is involved and partly responsible for the memory disturbances seen in aged animals and humans. Nicotine receptor modulator may therefore be useful in the

treatment of Alzheimer's disease, memory loss, memory dysfunction, AIDS-dementia, senile dementia or neurodegenerative disorders.

Parkinsons disease appears to involve degeneration of dopaminergic neurons. One symptom of the disease has been observed to be loss of nicotinic receptors associated with the

- 5 dopaminergic neurons and possibly interfering with the process of release of dopamine. As sustained nicotine administration increases the number of receptors present, administration of nicotine receptor modulators may ameliorate the symptoms of Parkinson's disease. Other condition or disorders or disease ascribed to deficiencies in the dopaminergic system is: drug addiction, depression, obesity and narcolepsy.

- 10 Tourette's syndrome is a neuropsychiatric disorder involving a range of neurological and behavioral symptoms. It is believed that neurotransmitter dysfunction is involved though the pathophysiology is still unknown and that nicotine will be beneficial in the treatment of the disease (Devor et. al. The Lancet, vol. 8670 p. 1046, 1989)

- Schizophrenia is a severe psychiatric illness. Neuroleptic compounds has been used in the
15 treatment of the disease, the effect of the compounds is believed to be interaction in the dopaminergic system. Nicotine is proposed to be effective in the treatment of schizophrenia (ie. Adler et. al. Biol. Psychiatry, Vol. 32, p. 607-616, 1992.)

- Nicotine has been reported to have an effect on neurotransmitter release in several systems. Release of acetylcholine and dopamine by neurons upon administration of nicotine has been
20 reported (J. Neurochem. vol. 43, 1593-1598, 1984) and release of norepinephrine by Hall et. al. (Biochem. Pharmacol. vol. 21, 1829-1838, 1972) Release of serotonin by Hery et. al. (Arch. Int. Pharmacodyn. Ther. vol. 296. p. 91-97, 1977). Release of glutamate by Toth et. al (Neurochem. Res. vol. 17, p. 265-271, 1992)

- The serotonin system and dysfunction's of the serotonergic system is believed to be involved
25 in diseases or conditions or disorders like: anxiety, depression, eating disorders, obsessive compulsive disorder, panic disorders, chemical substance abuse, alcoholism, pain, memory deficits and anxiety, pseudodementia, Ganster's syndrome, migraine pain, bulimia, obesity, pre-menstrual syndrome or late luteal phase syndrome, tobacco abuse, post-traumatic syndrome, social phobia, chronic fatigue syndrome, premature ejaculation, erectile difficulty,
30 anorexia nervosa, disorders of sleep, autism, mutism or trichotillomania.

Nicotine improves concentration and task performance. Therefore compounds exhibiting nicotine receptor modulating properties will be likely to be useful compounds in the treatment of learning deficit, cognition deficit, attention deficit, attention deficit hyperactivity disorder and dyslexia.

- 35 Tobacco use and especially cigarette smoking is recognised as a serious health problem. However nicotine withdrawal symptoms associated with smoking cessation makes it difficult to

break this habit. Withdrawal symptoms include anger, anxiety, difficulties in concentrating, restlessness, decreased heart rate and increased appetite and weight gain. Nicotine itself has shown to ease the withdrawal symptoms.

- 5 Withdrawal from addictive substances, i.e. opiates, benzodiazepines, ethanol, tobacco or nicotine, is in general a traumatic experience characterised by anxiety and frustration. Nicotine has been found to be effective in reducing anger, irritability, frustration and feelings of tension without causing general response depression, drowsiness or sedation and compounds having same characteristics as nicotine is likely to have same effects.

10

Mild to moderate pain is normally treatable with NSAID's (non-steroidal anti-inflammatory drugs) while opiates are used preferentially for moderate to severe pain. The opiates have some well-known side-effects, including chemical dependence and abuse potential as well as a depressive effect on the respiratory and gastrointestinal system. There exists therefore a strong need for analgesic compounds that do not exhibit these side effects and which can relieve mild, moderate and severe pain of acute, chronic or recurrent character as well as migraine pain and postoperative pain, phantom limb pain.

15

Epibatidine, a compound isolated from the skin of a poison frog, is a very potent analgesic with an approximate potency of 500 times that of morphine. The analgesic effect is not affected by naloxone, which is an indication of a negligible affinity for the opiate receptors. Epibatidine is an nicotinic cholinergic receptor agonist and it is therefore very likely, that compounds possessing this receptor modulating character will also show a strong analgesic response. The compounds of the present invention has proven useful for modulation of smooth muscle contractions and may therefore be used in the treatment or prevention of condition or disorders or diseases inherent from smooth muscle contractions like i.e. convulsive disorders, angina pectoris, premature labor, convulsions, diarrhoea, asthma, epilepsy, tardive dyskinesia, hyperkinesia.

20

25

Further, it is well known that nicotine has an effect on appetite and it is predicted that modulators at the nicotine ACh receptor may be useful as appetite suppressants in the treatment of obesity and eating disorders.

30

The cholinergic receptors play an important role in the functioning of muscles, organs and generally in the central or peripheral system. There are also complex interactions between cholinergic receptors and the function of receptors of other neurotransmitters such as dopamine, serotonin and noradrenaline.

35

It is likely that nicotine receptor modulator compounds can be effective in preventing or treating conditions or disorders or diseases like: inflammation, inflammatory skin conditions, Chron's disease, inflammatory bowel disease, irritable colon, ulcerative colitis, irritable colon, diarrhoea, neurodegeneration, peripheral neuropathy, amyotrophic lateral sclerosis, 5 nociception, endocrine disorders, thyrotoxicosis, pheochromocytoma, hypertension, arrhythmias, mania, manic depression, Huntington's disease, jetlag.

The compounds of the present invention are nicotine receptor modulators and has the potential to exhibit nicotinic pharmacology, preferentially without the side effects associated 10 with nicotine itself. Additionally, the compounds are expected to have the potential as enhancers of neurotransmitter secretion and suppress symptoms associated with a low activity of neurotransmitters.

Objects of the Invention

15

It is an object of the present invention to provide novel heteroaryl diazacycloalkane derivatives which are useful for the treatment of a range of diseases and disorders characterized by decreased cholinergic function or responsive to the activity of nicotinic ACh receptor agonists.

20 Another object of the present invention is to provide novel pharmaceutical compositions containing these compounds, as well as methods for the preparation thereof and methods for the treatment therewith.

It is yet another object of the invention to provide novel compounds that have some if not all of 25 the following favourable characteristics:

- A selective binding to the receptor subtypes of neuronal nAChR's, e.g. the non- $\alpha 7$ subtypes.
- A low affinity for the muscular subtype.
- A oral efficacy in an *in vivo* (rat model) of arousal/attention.
- 30 - A low toxicity *in vivo*.
- No adverse effects on heart rate or blood pressure *in vivo*.
- Compounds that are non-mutagenic.


Other objects will become apparent hereinafter to one skilled in the art.

35

Summary of the invention

The invention then, inter alia, comprises the following, alone or in combination:

- 5 A compound represented by the general formula



- any of its enantiomers or any mixture thereof, isotopes thereof or a pharmaceutically
 10 acceptable salt thereof;

wherein

- n is 1, 2 or 3;
 m is 0, 1 or 2;
 15 R represents hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, or aralkyl; and

R¹ represents **aminophenyl; nitrophenyl; hydroxyphenyl, alkoxyphenyl;**

- a monocyclic 5 to 6 membered heterocyclic group** which may be substituted one or more
 times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl,
 20 alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen,
 CF₃, OCF₃, CN, amino, nitro, -COOR³, -CONR²R³, -NH-CO₂R², NHCO-R², -OCO-NR²R³;
 wherein R² and R³ independently represents hydrogen or alkyl;

- aryl optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl
 alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃,
 25 CN, amino and nitro;

-X-alkyl-Y-alkyl wherein X and Y independently represents O, S, NH, N-alkyl or Se;
 and alkyl is optionally substituted with alkoxy or thioalkoxy;

- X-(alkyl)_o-aryl wherein o is 0 or 1 and X represents O, S, NH, N-alkyl or Se;
 optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl,
 30 alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and
 nitro;

-X-(alkyl)_o-Z wherein o is 0 or 1 and X represents O, S, NH, N-alkyl or Se and Z represents a 5- or 6- membered monocyclic heterocyclic group; optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

- 5 - a monocyclic 5 to 6 membered heterocyclic group optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

or R¹ represents a **bicyclic heterocyclic group**, composed of a 5 to 6 membered monocyclic
 10 heterocyclic group fused to a benzene ring, and which may be substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, alkoxy-alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino, nitro, aryl optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy,
 15 halogen, CF₃, OCF₃, CN, amino and nitro; and a monocyclic 5 to 6 membered heterocyclic group optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

- 20 with the proviso that when R¹ is hydroxyphenyl, alkoxyphenyl, nitrophenyl or aminophenyl, then m and n are not both 2;

A pharmaceutical composition, comprising a therapeutically effective amount of a compound of above, or a pharmaceutically acceptable addition salt thereof, together with at least one
 25 pharmaceutically acceptable carrier or diluent.

The use of a compound as above for the manufacture of a medicament for the treatment of a disease of a living animal body, including a human, which disease is responsive to the activity of nicotinic ACh receptor modulators.

30

A compound as above for use as a medicament for treatment of a disease of a living animal body, including a human, which disease is responsive to the activity of nicotinic Ach receptors modulators;

A method of treating a disease of a living animal body, including a human, which disease is responsive to the activity of nicotinic ACh receptor modulators, comprising the step of administering to such a living animal body, including a human, in need thereof a therapeutically effective amount of a compound as above;

5

Detailed disclosure of the invention

A preferred aspect of the invention comprises a compound of formula (I) as above wherein the group bridging R and R¹ is a piperazine, a homopiperazine, a 1,4-di-azacyclooctane, a 1,5-di-azacyclooctane, a 1,3-diazacyclohexane or an imidazolidine;

Another preferred aspect of the invention comprises compounds of formula (I) as above wherein R is as defined above and R¹ represents an optionally substituted heteroaryl attached to a piperazine, a homopiperazine, a 1,4-diazacyclooctane, a 1,5-diazacyclooctane, an imidazolidine or a 1,3-diazacyclohexane;

Another preferred aspect of the invention comprises compound of formula (I), as above wherein R¹ represents **isoquinolinyl**; or **phenyl** optionally substituted with nitro, hydroxy, amino or alkoxy; or **pyridyl, quinolinyl, pyridazinyl or pyridinyl** all of which may be optionally substituted one or more times with alkoxy, cycloalkoxy, alkoxyalkoxy, alkoxycycloalkyl, hydroxyalkoxy, thioalkoxy, thioalkylaryl, alkenoxy, alkynoxy, carboxylamido, a bicyclic heterocyclic group, thioalkoxyaryl, thioaryl, hydroxy or trifluoromethanesulfonyloxy; halogen, phenyl optionally substituted with nitro; a monocyclic 5 to 6 membered heterocyclic group optionally substituted with alkyl;

25

Yet another aspect of the invention comprises a compound of formula (I) as above, wherein R¹ represents 3-pyridyl, 3-quinolinyl, 5-methoxy-3-pyridyl, 5-chloro-3-pyridyl, 5-(4-methyl-1-piperazinyl)-3-pyridyl, 5-ethoxy-3-pyridyl, 6-methoxy-3-pyridyl, 5-propyloxy-3-pyridinyl, 5-phenyl-3-pyridyl, 5-(2-methyl-propyloxy)-3-pyridyl, 5-propyl-1-en-oxy-3-pyridyl, 2-chloro-5-methoxy-3-pyridyl, 6-chloro-5-methoxy-3-pyridyl, 6-bromo-5-ethoxy-3-pyridyl, 6-(N-pyrrolidinyl)-3-pyridyl, 6-phenyl-3-pyridyl, 5-(3-nitrophenyl)-3-pyridyl, 5-butoxy-3-pyridyl, 5-methoxyethoxy-3-pyridyl, 5-(2-methyl-propoxy)-3-pyridyl, 5-(2-hydroxy-ethoxy)-3-pyridyl, 5-(3-methyl-butoxy)-3-pyridyl, 5-cyclopropylmethoxy-3-pyridyl, 5-propyloxy-3-pyridyl, 5-hexyloxy-3-pyridyl, 5-cyclohexylmethoxy-3-pyridyl, 6-thioethoxy-3-pyridyl, 5-(2-ethoxy-ethoxy)-3-pyridyl, 5-pentyloxy-3-pyridyl, 5-heptyloxy-3-pyridyl, 5-(propyl-1-en-oxy)-3-pyridyl, 5-thiobenzyl-3-pyridyl,

35

5-carboxylamido-3-pyridyl, 5-thiophenyl-3-pyridyl, 5-methoxy-methoxy-3-pyridyl, 5-(3-pyridyl)-3-pyridyl, 5-(1-pyrrolyl)-3-pyridyl, 5-(1-indolyl)-3-pyridyl, 5,6-dimethoxy-3-pyridyl, 5-ethenyloxy-3-pyridyl, 5-cyclopentyloxy-3-pyridyl, 5-(ethoxy-d5)-3-pyridyl, 3-chloro-5-pyridyl, 3-bromo-5-pyridyl, 4-isoquinolyl, 5-hydroxy-3-pyridyl, 5-trifluoromethanesulfonyl-oxy-3-pyridyl, 5-ethynyl-3-pyridyl, 3-nitrophenyl, 3-aminophenyl, 3-methoxyphenyl, 3-hydroxyphenyl, 6-chloro-3-pyridazinyl, 6-phenyl-3-pyridazinyl, 6-chloro-2-pyrazinyl, 3,6-dimethyl-2-pyrazinyl, 6-methyl-3-pyridazinyl, 5-trifluoromethyl-3-pyridyl, 6-bromo-3-pyridyl, 6-chloro-3-pyridyl or 3-pyridazinyl;

A preferred aspect of the invention comprises a compound of formula (I) as above wherein R
10 represents hydrogen, lower alkyl or aralkyl;

The most preferred aspect of the invention comprises a compound of formula (I), as above, said compound being:

- 15 4-Methyl-1-(3-pyridyl)-piperazine;
- 4-Methyl-1-(3-pyridyl)-piperazine;
- 4-Methyl-1-(5-methoxy-3-pyridyl)-piperazine;
- 3,5-Bis-[4,4'-methyl-1,1'-piperazinyl]pyridine;
- 1-(5-Chloro-3-pyridyl)-4-Methyl-piperazine;
- 20 4-Methyl-1-(5-phenyl-3-pyridyl)-piperazine;
- 1-(5-Ethoxy-3-pyridyl)-4-methyl-piperazine;
- 1-(5-Butoxy-3-pyridyl)-4-methyl-piperazine;
- 4-Methyl-1-[5-(propyl-1-en-oxy)-3-pyridyl]-piperazine;
- 1-(5-Ethenyloxy-3-pyridyl)-4-methyl-piperazine;
- 25 1-(5-Methoxy-3-pyridyl)-4-methyl-1,5-diazacyclooctane;
- 1-(6-Chloro-3-pyridazinyl)-4-methyl-1,5-diazacyclooctane;
- 1-(3-Pyridyl)-piperazine;
- 1-(3-Pyridyl)-piperazine;
- 1-(5-Methoxy-3-pyridyl)-piperazine;
- 30 1-(5-Chloro-3-pyridyl)-piperazine;
- 1-(5-Phenyl-3-pyridyl)-piperazine;
- 1-(5-Methoxy-3-pyridyl)-1,5-diazacyclooctane;
- 1-(6-Chloro-3-pyridazinyl)-1,5-diazacyclooctane;
- 1-(6-Chloro-3-pyridazinyl)-1,4-diazacyclooctane;
- 35 1-(5-Ethoxy-3-pyridyl)piperazine;

- 1-(5-Butoxy-3-pyridyl)piperazine;
1-[5-(Propyl-1-en-oxy)-3-pyridyl]-piperazine;
1-(5-Ethenyloxy-3-pyridyl)piperazine;
1-(5-Ethoxy-3-pyridyl)-1,5-diazacyclooctane,
5 1-(5-Propyloxy-3-pyridinyl)-1,5-diazacyclooctane,
1-[5-(Propyl-1-en-oxy)-3-pyridyl]-1,5-diazacyclooctane,
1-(5-Ethenyloxy-3-pyridyl) 1,5-diazacyclooctane;
1-(5-Ethoxy-3-pyridyl)-4-ethyl-piperazine;
4-Methyl-1-(3-pyridyl)-homopiperazine;
10 4-Methyl-1-(3-pyridyl)-homopiperazine;
1-(5-Methoxy-3-pyridyl)-4-methyl-homopiperazine;
1-(5-Ethoxy-3-pyridyl)-4-methyl-homopiperazine;
4-Methyl-1-(5-phenyl-3-pyridyl)-homopiperazine;
1-(5-Butoxy-3-pyridyl)-4-methyl-homopiperazine;
15 1-(5-Methoxyethoxy-3-pyridyl)-4-methyl-homopiperazine;
4-Methyl-1-[5-(2-methyl-propyloxy)-3-pyridyl]-homopiperazine;
1-(5-Cyclopropylmethoxy-3-pyridyl)-4-methyl-homopiperazine;
4-Methyl-1-(5-propyloxy-3-pyridyl)-homopiperazine;
1-(5-Hexyloxy-3-pyridyl)-4-methyl-homopiperazine;
20 4-Methyl-1-[5-(3-methyl-butoxy)-3-pyridyl]-homopiperazine;
4-Methyl-1-(6-thioethoxy-3-pyridyl)-homopiperazine;
1-(5-Cyclohexylmethoxy-3-pyridyl)-4-methyl-homopiperazine;
4-Methyl-1-(5-pentyloxy-3-pyridyl)-homopiperazine;
1-(5-Heptyloxy-3-pyridyl)-4-methyl-homopiperazine;
25 4-Methyl-1-(5-propyl-1-en-oxy-3-pyridyl)-homopiperazine;
4-Methyl-1-(5-thiobenzyl-3-pyridyl)-homopiperazine;
4-Methyl-1-[5-(3-pyridyl)-3-pyridyl]-homopiperazine;
4-Methyl-1-(3-nitrophenyl)-homopiperazine;
4-Methyl-1-(6-chloro-3-pyridazinyl)-homopiperazine;
30 4-Methyl-1-(6-phenyl-3-pyridazinyl)-homopiperazine;
4-Methyl-1-(3-pyridazinyl)-homopiperazine;
4-Methyl-1-(6-methyl-3-pyridazinyl)-homopiperazine;
1-(5-Cyclopentyloxy-3-pyridyl)-4-methyl-homopiperazine;
4-Benzyl-1-(3-pyridyl)-homopiperazine;
35 4-Ethyl-1-(3-pyridyl)-homopiperazine;

- 1-(3-Pyridyl)-homopiperazine;
1-(3-Pyridyl)-homopiperazine;
1-(6-Methoxy-3-pyridyl)-homopiperazine;
1-(2-Chloro-5-methoxy-3-pyridyl)-homopiperazine;
5 1-(6-Chloro-5-methoxy-3-pyridyl)-homopiperazine;
1-(6-Bromo-5-ethoxy-3-pyridyl)-homopiperazine;
1-[6-(N-Pyrrolidiny)-3-pyridyl]-homopiperazine;
1-(6-Phenyl-3-pyridyl)-4-homopiperazine;
1-[5-(3-Nitrophenyl)-3-pyridyl]-4-homopiperazine;
10 1-(5-Methoxy-3-pyridyl)-homopiperazine;
1-(5-Phenyl-3-pyridyl)-homopiperazine;
1-(5-Ethoxy-3-pyridyl)-homopiperazine;
1-(5-Butoxy-3-pyridyl)-homopiperazine;
1-(5-Methoxyethoxy-3-pyridyl)-homopiperazine;
15 1-[5-(2-Methyl-propoxy)-3-pyridyl]-homopiperazine;
1-[5-(2-Hydroxy-ethoxy)-3-pyridyl]-homopiperazine;
1-[5-(3-Methyl-butoxy)-3-pyridyl]-homopiperazine;
1-(5-Cyclopropylmethoxy-3-pyridyl)-homopiperazine;
1-(5-Propyloxy-3-pyridyl)-homopiperazine;
20 1-(5-Hexyloxy-3-pyridyl)-homopiperazine;
1-(5-Cyclohexylmethoxy-3-pyridyl)-homopiperazine;
1-(6-Thioethoxy-3-pyridyl)-homopiperazine;
1-[5-(2-Ethoxy-ethoxy)-3-pyridyl]-homopiperazine;
1-(5-Pentyloxy-3-pyridyl)-homopiperazine;
25 1-(5-Heptyloxy-3-pyridyl)-homopiperazine;
1-[5-(Propyl-1-en-oxy)-3-pyridyl]-homopiperazine;
1-(5-Thiobenzyl-3-pyridyl)-homopiperazine;
1-(5-Carboxylamido-3-pyridyl)-homopiperazine;
1-(5-Thiophenyl-3-pyridyl)-homopiperazine;
30 1-[(5-Methoxy-methoxy)-3-pyridyl]-homopiperazine;
1-[5-(3-Pyridyl)-3-pyridyl]-homopiperazine;
1-(5-(1-Pyrrolyl)-3-pyridyl)-homopiperazine;
1-(5-(1-Indolyl)-3-pyridyl)-homopiperazine;
1-(5,6-Dimethoxy-3-pyridyl)-homopiperazine;
35 1-(5-Ethenyloxy-3-pyridyl)homopiperazine;

- 1-(5-Cyclopentyloxy-3-pyridyl)homopiperazine;
- 1-[5-(Ethoxy-d5)-3-pyridyl]-homopiperazine;
- 1-(3-Chloro-5-pyridyl)homopiperazine;
- 1-(3-Bromo-5-pyridyl)homopiperazine;
- 5 1-(4-Isoquinoliny)-homopiperazine;
- 1-(5-Hydroxy-3-pyridyl)-homopiperazine;
- 1-(5-Trifluoromethanesulfonyl-oxy-3-pyridyl)-homopiperazine;
- 1-(5-Ethynyl-3-pyridyl)-homopiperazine;
- 1-(3-Nitrophenyl)-homopiperazine;
- 10 1-(3-Aminophenyl)-homopiperazine;
- 1-(3-Methoxyphenyl)-homopiperazine;
- 1-(3-Hydroxyphenyl)-homopiperazine;
- 1-(6-Chloro-3-pyridazinyl)-homopiperazine;
- 1-(6-Phenyl-3-pyridazinyl)-homopiperazine;
- 15 1-(6-Chloro-2-pyrazinyl)-homopiperazine;
- 1-(3,6-Dimethyl-2-pyrazinyl)-homopiperazine;
- 1-(6-Methyl-3-pyridazinyl)-homopiperazine;
- 1-(5-Trifluoromethyl-3-pyridyl)-homopiperazine;
- 1-(6-Bromo-3-pyridyl)-homopiperazine;
- 20 1-(6-Chloro-3-pyridyl)-homopiperazine;
- 1-(3-Pyridazinyl)-homopiperazine;
- N-(5-Methoxy-3-pyridyl)-ethylenediamine;
- 1-(5-Methoxy-3-pyridyl)-imidazolidine;
- 1-(5-Methoxy-3-pyridyl)-1.3-diazacyclohexane;
- 25 1-(5-Benzyloxy-3-pyridyl)-piperazine;
- 1-(5-Benzyloxy-3-pyridyl)-homopiperazine;
- 4-Methyl-1-(5-benzyloxy-3-pyridyl)-piperazine;
- 4-Methyl-1-(5-benzyloxy-3-pyridyl)-homopiperazine;
- 1-(5-Thiobenzyloxy-3-pyridyl)-piperazine;
- 30 4-Methyl-1-(5-thiobenzyloxy-3-pyridyl)-piperazine;
- 1-(5-Phenoxy-3-pyridyl)-piperazine;
- 1-(5-Phenoxy-3-pyridyl)-homopiperazine;
- 4-Methyl-1-(5-phenoxy-3-pyridyl)-piperazine;
- 4-Methyl-1-(5-phenoxy-3-pyridyl)-homopiperazine;
- 35 1-(5-Thiophenoxy-3-pyridyl)-piperazine;

- 4-Methyl-1-(5-thiophenoxy-3-pyridyl)-piperazine;
4-Methyl-1-(5-thiophenoxy-3-pyridyl)-homopiperazine;
1-(5-(3-Pyridyloxy)-3-pyridyl)-piperazine;
1-(5-(3-Pyridyloxy)-3-pyridyl)-homopiperazine;
5 4-Methyl-1-(5-(3-pyridyloxy)-3-pyridyl)-piperazine;
4-Methyl-1-(5-(3-pyridyloxy)-3-pyridyl)-homopiperazine;
1-(5-(2-Thiopyridyloxy)-3-pyridyl)-piperazine;
1-(5-(2-Thiopyridyloxy)-3-pyridyl)-homopiperazine;
4-Methyl-1-(5-(2-thiopyridyloxy)-3-pyridyl)-piperazine;
10 4-Methyl-1-(5-(2-thiopyridyloxy)-3-pyridyl)-homopiperazine;
1-(5-(4-Thiopyridyloxy)-3-pyridyl)-piperazine;
1-(5-(4-Thiopyridyloxy)-3-pyridyl)-homopiperazine;
4-Methyl-1-(5-(4-thiopyridyloxy)-3-pyridyl)-piperazine;
4-Methyl-1-(5-(4-thiopyridyloxy)-3-pyridyl)-homopiperazine;
15 1-(5-(2-Pyridyl)-3-pyridyl)-piperazine;
1-(5-(3-Pyridyl)-3-pyridyl)-piperazine;
1-(5-(4-Pyridyl)-3-pyridyl)-piperazine;
1-(5-(2-Pyridyl)-3-pyridyl)-homopiperazine;
1-(5-(4-Pyridyl)-3-pyridyl)-homopiperazine;
20 4-Methyl-1-(5-(2-pyridyl)-3-pyridyl)-piperazine;
4-Methyl-1-(5-(3-pyridyl)-3-pyridyl)-piperazine;
4-Methyl-1-(5-(4-pyridyl)-3-pyridyl)-piperazine;
4-Methyl-1-(5-(2-pyridyl)-3-pyridyl)-homopiperazine;
4-Methyl-1-(5-(4-pyridyl)-3-pyridyl)-homopiperazine;
25 1-(5-(2-Furanyl)-3-pyridyl)piperazine;
4-Methyl-1-(5-(2-furanyl)-3-pyridyl)piperazine;
1-(5-(3-Furanyl)-3-pyridyl)piperazine;
4-Methyl-1-(5-(3-furanyl)-3-pyridyl)piperazine;
1-(5-(2-Thienyl)-3-pyridyl)piperazine;
30 1-(5-(2-Thienyl)-3-pyridyl)homopiperazine;
4-Methyl-1-(5-(2-thienyl)-3-pyridyl)piperazine;
4-Methyl-1-(5-(2-thienyl)-3-pyridyl)homopiperazine;
1-(5-(3-Thienyl)-3-pyridyl)piperazine;
1-(5-(3-Thienyl)-3-pyridyl)homopiperazine;
35 4-Methyl-1-(5-(3-thienyl)-3-pyridyl)piperazine;

- 4-Methyl-1-(5-(3-thienyl)-3-pyridyl)homopiperazine;
1-(5-(2-Thiazolyl)-3-pyridyl)piperazine;
1-(5-(2-Thiazolyl)-3-pyridyl)homopiperazine;
4-Methyl-1-(5-(2-thiazolyl)-3-pyridyl)piperazine;
5 4-Methyl-1-(5-(2-thiazolyl)-3-pyridyl)homopiperazine;
1-(5-(Methyl-ethylendioxy)-3-pyridyl)piperazine;
4-Methyl-1-(5-(methyl-ethylendioxy)-3-pyridyl)piperazine;
1-(5-(Ethyl-ethylendioxy)-3-pyridyl)piperazine;
4-Methyl-1-(5-(ethyl-ethylendioxy)-3-pyridyl)piperazine;
10 1-(5-(Butyl-ethylendioxy)-3-pyridyl)piperazine;
1-(5-(Butyl-ethylendioxy)-3-pyridyl)homopiperazine;
4-Methyl-1-(5-(butyl-ethylendioxy)-3-pyridyl)piperazine;
4-Methyl-1-(5-(butyl-ethylendioxy)-3-pyridyl)homopiperazine;
1-(5-(Propyl-ethylendioxy)-3-pyridyl)piperazine;
15 1-(5-(Propyl-ethylendioxy)-3-pyridyl)homopiperazine;
4-Methyl-1-(5-(propyl-ethylendioxy)-3-pyridyl)piperazine;
4-Methyl-1-(5-(propyl-ethylendioxy)-3-pyridyl)homopiperazine;
1-(5-(1,4,7-Trioxanonyl)-3-pyridyl)piperazine;
1-(5-(1,4,7-Trioxanonyl)-3-pyridyl)homopiperazine;
20 4-Methyl-1-(5-(1,4,7-trioxanonyl)-3-pyridyl)piperazine;
4-Methyl-1-(5-(1,4,7-trioxanonyl)-3-pyridyl)homopiperazine;
1-(5-(1,7-Dioxa-4-thia-octanyl)-3-pyridyl)piperazine;
1-(5-(1,7-Dioxa-4-thia-octanyl)-3-pyridyl)homopiperazine;
4-Methyl-1-(5-(1,7-dioxa-4-thia-octanyl)-3-pyridyl)piperazine;
25 4-Methyl-1-(5-(1,7-dioxa-4-thia-octanyl)-3-pyridyl)homopiperazine;
1-(5-(2-Methylthio-ethoxy)-3-pyridyl)piperazine;
1-(5-(2-Methylthio-ethoxy)-3-pyridyl)homopiperazine;
4-Methyl-1-(5-(2-methylthio-ethoxy)-3-pyridyl)piperazine;
4-Methyl-1-(5-(2-methylthio-ethoxy)-3-pyridyl)homopiperazine;
30 1-(5-(2-Ethylthio-ethoxy)-3-pyridyl)piperazine;
1-(5-(2-Ethylthio-ethoxy)-3-pyridyl)homopiperazine;
4-Methyl-1-(5-(2-ethylthio-ethoxy)-3-pyridyl)piperazine;
4-Methyl-1-(5-(2-ethylthio-ethoxy)-3-pyridyl)homopiperazine;
1-(5-(Cyclopropylmethoxy)-3-pyridyl)piperazine;
35 4-Methyl-1-(5-(cyclopropylmethoxy)-3-pyridyl)piperazine;

- 1-(5-(sec-Butoxy)-3-pyridyl)piperazine;
 4-Methyl-1-(5-(sec-butoxy)-3-pyridyl)piperazine;
 1-(5-(Isopentoxy)-3-pyridyl)piperazine;
 4-Methyl-1-(5-(isopentoxy)-3-pyridyl)piperazine;
 5 1-(5-(Isopropoxy)-3-pyridyl)piperazine;
 1-(5-(Isopropoxy)-3-pyridyl)homopiperazine;
 4-Methyl-1-(5-(isopropoxy)-3-pyridyl)piperazine;
 4-Methyl-1-(3-pyridyl)-1,4-diazacyclooctane;
 1-(3-Pyridyl)-1,4-diazacyclooctane;
 10 1-(3-Pyridyl)-1,5-diazacyclooctane
 4-Methyl-1-(3-pyridyl)-1,5-diazacyclooctane;
- or a pharmaceutically acceptable addition salt thereof;

15 **Pharmaceutical acceptable addition salts**

The chemical compound of the invention may be provided in any form suitable for the intended administration. Suitable forms include pharmaceutically (i.e. physiologically) acceptable salts, and pre- or prodrug forms of the chemical compound of the invention.

- 20 Examples of pharmaceutically acceptable addition salts include, without limitation, the non-toxic inorganic and organic acid addition salts such as the acetate derived from acetic acid, the aconate derived from aconitic acid, the ascorbate derived from ascorbic acid, the benzenesulfonate derived from benzenesulfonic acid, the benzoate derived from benzoic acid, the cinnamate derived from cinnamic acid, the citrate derived from citric acid, the embonate
 25 derived from embonic acid, the enantate derived from enanthic acid, the formate derived from formic acid, the fumarate derived from fumaric acid, the glutamate derived from glutamic acid, the glycolate derived from glycolic acid, the hydrochloride derived from hydrochloric acid, the hydrobromide derived from hydrobromic acid, the lactate derived from lactic acid, the maleate derived from maleic acid, the malonate derived from malonic acid, the mandelate derived from
 30 mandelic acid, the methanesulfonate derived from methane sulphonic acid, the naphthalene-2-sulphonate derived from naphthalene-2-sulphonic acid, the nitrate derived from nitric acid, the perchlorate derived from perchloric acid, the phosphate derived from phosphoric acid, the phthalate derived from phthalic acid, the salicylate derived from salicylic acid, the sorbate derived from sorbic acid, the stearate derived from stearic acid, the succinate derived from
 35 succinic acid, the sulphate derived from sulphuric acid, the tartrate derived from tartaric acid,

the toluene-p-sulphonate derived from p-toluene sulfonic acid, and the like. Such salts may be formed by procedures well known and described in the art.

Other acids such as oxalic acid, which may not be considered pharmaceutically acceptable, may be useful in the preparation of salts useful as intermediates in obtaining a chemical

5 compound of the invention and its pharmaceutically acceptable acid addition salt.

Metal salts of a chemical compound of the invention includes alkali metal salts, such as the sodium salt of a chemical compound of the invention containing a carboxy group.

The chemical compound of the invention may be provided in solved or unsolved form together with a pharmaceutically acceptable solvents such as water, ethanol and the like. In general,

10 solved forms are considered equivalent to dissolved forms for the purposes of this invention.

Steric isomers

The chemical compounds of the present invention may exist in (+) and (-) forms as well as in
15 racemic forms. The racemates of these isomers and the individual isomers themselves are within the scope of the present invention.

Racemic forms can be resolved into the optical antipodes by known methods and techniques.

One way of separating the diastereomeric salts is by use of an optically active acid, and liberating the optically active amine compound by treatment with a base. Another method for
20 resolving racemates into the optical antipodes is based upon chromatography on an optical active matrix. Racemic compounds of the present invention can thus be resolved into their optical antipodes, e.g., by fractional crystallisation of d- or l- (tartrates, mandelates, or camphorsulphonate) salts for example.

The chemical compounds of the present invention may also be resolved by the formation of
25 diastereomeric amides by reaction of the chemical compounds of the present invention with an optically active activated carboxylic acid such as that derived from (+) or (-) phenylalanine, (+) or (-) phenylglycine, (+) or (-) camphanic acid or by the formation of diastereomeric carbamates by reaction of the chemical compound of the present invention with an optically active chloroformate or the like.

30 Additional methods for the resolving the optical isomers are known in the art. Such methods include those described by *Jaques J, Collet A, & Wilen S* in "Enantiomers, Racemates, and Resolutions", John Wiley and Sons, New York (1981).

Moreover, some of the chemical compounds of the invention contains double bonds and may thus exist in two forms, cis- and trans-forms (Z- and E-form), depending on the arrangement

of the substituents around the -C=C- double bond. A chemical compound of the present invention may thus be the cis- or trans-form (Z- and E-form), or it may be a mixture hereof.

Definition of substituents

5

Halogen is fluorine, chlorine, bromine or iodine.

Alkyl means a straight chain or branched chain of one to eight carbon atoms, including but not limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, pentyl, and hexyl;

10

Cycloalkyl means cyclic alkyl of three to seven carbon atoms, including but not limited to cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl;

Alkenyl means a group of from two to six carbon atoms, including at least one double bond, 15 for example, but not limited to ethenyl, 1,2- or 2,3-propenyl, 1,2-, 2,3-, or 3,4-butenyl.

Alkynyl means a group of from two to six carbon atoms, including at least one triple bond, for example, but not limited to ethynyl, 1,2- or 2,3-propynyl, 1,2- or 2,3- or 3,4-butyne.

20 Cycloalkylalkyl means cycloalkyl as above and alkyl as above, meaning for example, cyclopropylmethyl.

Alkoxy is O-alkyl, wherein alkyl is as defined above.

25 Cycloalkoxy is O-cycloalkyl, wherein cycloalkyl is as defined above.

Alkenoxy is O-alkenyl wherein alkenyl is as defined above.

Alkynoxy is O-alkynyl, wherein alkynyl is as defined above;

30

Thioalkoxy is S-alkyl, wherein alkyl is as defined above.

Amino is NH_2 or NH-alkyl or N-(alkyl)₂, wherein alkyl is as defined above.

A monocyclic 5- to 6-membered heterocyclic group includes, for example, oxazol-2-yl, oxazol-4-yl, oxazol-5-yl, isoxazol-3-yl, isoxazol-4-yl, isoxazol-5-yl, thiazol-2-yl, thiazol-4-yl, thiazol-5-yl, isothiazol-3-yl, isothiazol-4-yl, isothiazol-5-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl, 1,2,5-oxadiazol-3-yl, 1,2,5-oxadiazol-4-yl, 1,2,5-thiadiazol-3-yl, 1,2,5-thiadiazol-4-yl, 1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 2-furanyl, 3-furanyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrazinyl and 3-pyrazinyl and 1-pyrazolyl, 3-pyrazolyl, and 4-pyrazolyl.

- 10 A bicyclic heterocyclic group composed of a 5 to 6 membered monocyclic heterocyclic group and a fused benzene ring means a monocyclic 5 to 6 membered heterocyclic group as above which is fused to a benzene ring including, for example, 2-, 3-, 4-, 5-, 6-, 7-benzofuranyl, 1-, 2-, 4-, 5- benzimidazolyl, 2-, 3-, 4-, 5-, 6-, 7-indolyl, 2-,3-,4-,5-,6-,7-,8-quinoliny and 1-,3-,4-,5-,6-,7-,8-isoquinoliny.

15

Aryl is an aromatic hydrocarbon, such as phenyl and naphthyl.

Aralkyl means alkyl as above and aryl as above, meaning for example benzyl, phenethyl.

- 20 Isotopes means one or more atom in the compound is substituted with an isotope of the naturally occurring atoms and includes though not limited to deuterium, tritium, ^{13}C , ^{14}C , ^{131}I , ^{125}I , ^{123}I , ^{18}F ;

Further, the compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of this invention.

It will be appreciated by those skilled in the art that the compounds of the present invention may contain several chiral centers and that such compounds exist in the form of isomers (i.e. enantiomers). The invention includes all such isomers and any mixtures thereof including racemic mixtures.

Racemic forms can be resolved into the optical antipodes by known methods, for example, by separation of diastereomeric salts thereof with an optically active acid, and liberating the optically active amine compound by treatment with a base. Another method for resolving

racemates into the optical antipodes is based upon chromatography on an optically active matrix. Racemic compounds of the present invention can thus be resolved into their optical antipodes, e.g., by fractional crystallization of d- or l- (tartrates, mandelates, or camphorsulphonate) salts for example. The compounds of the present invention may also be
5 resolved by the formation of diastereomeric amides by reaction of the compounds of the present invention with an optically active activated carboxylic acid such as that derived from (+) or (-) phenylalanine, (+) or (-) phenylglycine, (+) or (-) camphanic acid or by the formation of diastereomeric carbamates by reaction of the compounds of the present invention with an optically active chloroformate or the like.

10

Additional methods for the resolution of optical isomers, known to those skilled in the art may be used, and will be apparent to the average worker skilled in the art. Such methods include those discussed by J. Jaques, A. Collet, and S. Wilen in "Enantiomers, Racemates, and Resolutions", John Wiley and Sons, New York (1981).

15

Optical active compounds can also be prepared from optical active starting materials.

The compounds of the invention may be prepared by any conventional method useful for the preparation of analogous compounds and as described in the examples below.

20

Starting materials for the processes described in the present patent application are known or can be prepared by known processes from commercially available materials

A compound of the invention can be converted to another compound of the invention using

25 conventional methods.

The products of the reactions described herein are isolated by conventional means such as extraction, crystallization, distillation, chromatography, and the like.

30

Biology

Nicotinic ACh receptors in the brain are pentameric structures composed of subunits distinct from those found in skeletal muscles. The existence of seven α -subunits ($\alpha 2$ - $\alpha 7$, $\alpha 9$) and three β -subunits ($\beta 2$ - $\beta 4$) in the mammalian brain has been described.

35

The predominant subtype with high affinity for nicotine is comprised of α_4 and β_2 subunits.

The affinity of compounds of the invention for nicotinic ACh receptors have been investigated in three test for *in vitro* inhibition of ^3H -epibatidin binding, ^3H - α -bunga-

5 rotoxin binding and ^3H -cytisine binding as described below:

***In vitro* inhibition of ^3H -cytisine binding**

The predominant subtype with high affinity for nicotine is comprised of α_4 and β_2 subunits.

10 nAChRs of the latter type can selectively be labelled by the nicotine agonist ^3H -cytisine.

Tissue Preparation: Preparations are performed at 0-4°C unless otherwise indicated. Cerebral cortices from male Wistar rats (150-250 g) are homogenized for 20 sec in 15 ml Tris, HCl (50 mM, pH 7.4) containing 120 mM NaCl, 5 mM KCl, 1 mM MgCl_2 and 2.5 mM CaCl_2 using an
15 Ultra-Turrax homogenizer. The homogenate is centrifuged at 27,000 x g for 10 min. The supernatant is discarded and the pellet is resuspended in fresh buffer and centrifuged a second time. The final pellet is resuspended in fresh buffer (35 ml per g of original tissue) and used for binding assays.

20 Assay: Aliquots of 500 μl homogenate are added to 25 μl of test solution and 25 μl of ^3H -cytisine (1 nM, final concentration), mixed and incubated for 90 min at 2°C. Non-specific binding is determined using (-)-nicotine (100 μM , final concentration). After incubation the samples are added 5 ml of ice-cold buffer and poured directly onto Whatman GF/C glass fibre filters under suction and immediately washed with 2 x 5 ml ice-cold buffer. The amount of
25 radioactivity on the filters is determined by conventional liquid scintillation counting. Specific binding is total binding minus non-specific binding.

***In vitro* inhibition of ^3H - α -bungarotoxin binding Rat brain**

30 α -Bungarotoxin is a peptide isolated from the venom of the Elapidae snake *Bungarus multicinctus* (Mebis et al., Biochem. Biophys. Res. Commun., 44(3), 711 (1971)) and has high affinity for neuronal and neuromuscular nicotinic receptors, where it acts as a potent antagonist. ^3H - α -Bungarotoxin binds to a single site in rat brain with an unique distribution pattern in rat brain (Clarke et al., J. Neurosci. 5, 1307-1315 (1985)).

^3H - α -Bungarotoxin labels nAChR formed by the α_7 subunit isoform found in brain and the α_1 isoform in the neuromuscular junction (Changeaux, Fidia Res. Found. Neurosci. Found. Lect. 4, 21-168 (1990). Functionally, the α_7 homo-oligomer expressed in oocytes has a calcium permeability greater than neuromuscular receptors and, in some instances greater than NMDA channels (Seguela et al., J. Neurosci. 13, 596-604 (1993).

Tissue preparation: Preparations are performed at 0-4°C unless otherwise indicated. Cerebral cortices from male Wistar rats (150-250 g) are homogenized for 10 sec in 15 ml 20 mM Hepes buffer containing 118 mM NaCl, 4.8 mM KCl, 1.2 mM MgSO_4 and 2.5 mM CaCl_2 (pH 7.5) using an Ultra-Turrax homogenizer. The tissue suspension is centrifuged at 27,000 x g for 10 min. The supernatant is discarded and the pellet is washed twice by centrifugation at 27,000 x g for 10 min in 20 ml fresh buffer, and the final pellet is resuspended in fresh buffer containing 0.01% BSA (35 ml per g of original tissue) and used for binding assays.

Assay: Aliquots of 500 μl homogenate are added to 25 μl of test solution and 25 μl of ^3H - α -bungarotoxin (2 nM, final concentration), mixed and incubated for 2 h at 37°C. Non-specific binding is determined using (-)-nicotine (1 mM, final concentration). After incubation the samples are added 5 ml of ice-cold Hepes buffer containing 0.05% PEI and poured directly onto Whatman GF/C glass fibre filters (presoaked in 0.1% PEI for at least 6 h) under suction and immediately washed with 2 x 5 ml ice-cold buffer. The amount of radioactivity on the filters is determined by conventional liquid scintillation counting. Specific binding is total binding minus non-specific binding.

***In vitro* inhibition of ^3H -epibatidin binding**

25

Epibatidin is an alkaloid that was first isolated from the skin of the Ecuadoran frog *Epipedobates tricolor* and was found to have very high affinity for neuronal nicotinic receptors, where it acts as a potent agonist. ^3H -epibatidin binds to two sites in rat brain, both of which have pharmacological profiles consistent with neuronal nicotinic receptors and a similar brain regional distribution (Hougling et al., Mol. Pharmacol. 48, 280-287 (1995)).

30

The high affinity binding site for ^3H -epibatidin is most certainly binding to the $\alpha_4\beta_2$ subtype of nicotinic receptors. The identity of the low affinity site is still unknown; does it represent a second nicotinic receptor or a second site in the same receptor. The inability of α -bungarotoxin

to compete for ^3H -epibatidin binding sites indicates that neither site measured represents the nicotinic receptor composed of α_7 subunits.

Tissue preparation: Preparations are performed at 0-4°C unless otherwise indicated. The

- 5 forebrain (\pm cerebellum) from a male Wistar rat (150-250 g) is homogenized for 10-20 sec in 20 ml Tris, HCl (50 mM, pH 7.4) using an Ultra-Turrax homogenizer. The tissue suspension is centrifuged at 27,000 x g for 10 min. The supernatant is discarded and the pellet is washed three times by centrifugation at 27,000 x g for 10 min in 20 ml fresh buffer, and the final pellet is resuspended in fresh buffer (400 ml per g of original tissue) and used for binding assays.
- 10 Assay: Aliquots of 2.0 ml homogenate are added to 0.100 ml of test solution and 0.100 ml of ^3H -epibatidin (0.3 nM, final concentration), mixed and incubated for 60 min at room temperature. Non-specific binding is determined using (-)-nicotine (30 μM , final concentration). After incubation the samples are poured directly onto Whatman GF/C glass fibre filters (pres soaked in 0.1% PEI for at least 20 min) under suction and immediately washed with 2 x 5
- 15 ml ice-cold buffer. The amount of radioactivity on the filters is determined by conventional liquid scintillation counting. Specific binding is total binding minus non-specific binding.

Compound	^3H - cytisine $\text{IC}_{50}(\mu\text{M})$	^3H - epibatidin $\text{IC}_{50}(\mu\text{M})$	^3H - α - bungarotoxin $\text{IC}_{50}(\mu\text{M})$
4-Methyl-1-(3-pyridyl)-piperazine fumaric acid salt (1A)	0.06	0.38	6.80
4-Methyl-1-(5-methoxy-3-pyridyl)-piperazine (3A)	0.068	0.38	32.50
1-(5-Chloro-3-pyridyl)-4-Methyl-piperazine fumaric acid salt (5A)	0.08	0.65	73.40
4-Methyl-1-(5-phenyl-3-pyridyl)-piperazine fumaric acid salt (6A)	0.08	1.10	>30
1-(5-Ethoxy-3-pyridyl)-4-methyl-piperazine fumaric acid salt (7A)	0.006	0.09	>30
1-(5-Methoxy-3-pyridyl)-4-methyl-1,5-diazacyclooctane fumaric acid salt (11A)	0.04	1.2	>30
1-(3-Pyridyl)-piperazine fumaric acid salt	0.13	0.63	>30

(1B)			
1-(5-Methoxy-3-pyridyl)-piperazine fumaric acid salt (3B)	0.80	4.50	>30
1-(5-Chloro-3-pyridyl)-piperazine fumaric acid salt (4B)	0.31	3.00	>30
1-(5-Methoxy-3-pyridyl)-1,5-diazacyclooctane fumaric acid salt (6B)	0.001	0.0032	2.5
1-(6-Chloro-3-pyridazinyl)-1,5-diazacyclooctane fumaric acid salt (7B)	0.03	0.15	0.46
1-(6-Chloro-3-pyridazinyl)-1,4-diazacyclooctane fumaric acid salt (8B)	0.55	0.92	5.50
1-(5-Ethoxy-3-pyridyl)piperazine fumaric acid salt (1D)	0.50	1.00	>30
1-(5-Butoxy-3-pyridyl)piperazine fumaric acid salt (2D)	0.18	1.40	>30
1-(5-Ethoxy-3-pyridyl)-4-ethyl-piperazine fumaric acid salt (9D)	0.07	0.21	>30
4-Methyl-1-(3-pyridyl)-homopiperazine (1E)	0.15	0.4	4.6
4-Methyl-1-(3-quinoliny)-homopiperazine fumaric acid salt (2E)	0.34	4.90	1.80
1-(5-Methoxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (3E)	0.094	0.75	26.00
1-(5-Ethoxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (4E)	0.05	0.30	>30
4-Methyl-1-(5-phenyl-3-pyridyl)-homopiperazine fumaric acid salt (5E)	0.20	1.50	16.0
1-(5-Butoxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (6E)	0.045	0.24	>30
1-(5-Methoxyethoxy-3-pyridyl)-4-methyl-homopiperazine (7E)	0.07	0.55	>30
4-Methyl-1-[5-(2-methyl-propyloxy)-3-pyridyl]-homopiperazine fumaric acid salt (8E)	0.07	0.45	>30

1-(5-Cyclopropylmethoxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (9E)	0.078	4.80	>30
4-Methyl-1-(5-propyloxy-3-pyridyl)-homopiperazine fumaric acid salt (10E)	0.06	0.29	>30
1-(5-Hexyloxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (11E)	0.80	0.25	>30
4-Methyl-1-[5-(3-methyl-butoxy)-3-pyridyl]-homopiperazine fumaric acid salt (12E)	0.055	0.27	>30
1-(5-Cyclohexylmethoxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (14E)	0.20	0.48	>30
4-Methyl-1-(5-pentyloxy-3-pyridyl)-homopiperazine fumaric acid salt (15E)	0.03	0.21	>30
1-(5-Heptyloxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (16E)	0.28	0.70	>30
<i>trans</i> -4-Methyl-1-(5-propyl-1-en-oxy-3-pyridyl)-homopiperazine fumaric acid salt (17E)	0.02	0.09	>30
4-Methyl-1-(5-thiobenzyl-3-pyridyl)-homopiperazine fumaric acid salt (18E)	0.60	0.80	15.0
4-Methyl-1-[5-(3-pyridyl)-3-pyridyl]-homopiperazine fumaric acid salt (19E)	0.17	1.80	>30
1-(5-Cyclopentyloxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (25E)	0.08	0.61	>30
1-(3-Pyridyl)-homopiperazine fumaric acid salt (1F)	0.002	0.005	0.19
1-(3-Quinoliny)-homopiperazine fumaric acid salt (2F)	0.006	0.084	5.90
1-(6-Chloro-5-methoxy-3-pyridyl)-homopiperazine fumaric acid salt (5F)	0.0007	0.005	0.90
1-(5-Methoxy-3-pyridyl)-homopiperazine fumaric acid salt (1H)	0.002	0.02	3.00

1-(5-Phenyl-3-pyridyl)-homopiperazine fumaric acid salt (2H)	0.003	0.02	37.00
1-(5-Ethoxy-3-pyridyl)-homopiperazine fumaric acid salt (3H)	0.002	0.01	20.00
1-(5-Methoxyethoxy-3-pyridyl)- homopiperazine fumaric acid salt (5H)	0.002	0.02	19.00
1-[5-(2-Methyl-propoxy)-3-pyridyl]- homopiperazine fumaric acid salt (6H)	0.002	0.01	>30
1-(5-Cyclopropylmethoxy-3-pyridyl)- homopiperazine fumaric acid salt (9H)	0.001	0.006	>30
1-(5-Propyloxy-3-pyridyl)-homopiperazine fumaric acid salt (10H)	0.0009	0.0054	>30
1-(6-Chloro-3-pyridazinyl)-homopiperazine fumaric acid salt (1K)	0.01	0.06	1.40
1-(6-Chloro-2-pyrazinyl)-homopiperazine fumaric acid salt (3K)	0.02	0.16	5.60
1-(6-Methyl-3-pyridazinyl)-homopiperazine fumaric acid salt (5K)	0.16	0.74	4.80
1-(5-Trifluoromethyl-3-pyridyl)- homopiperazine fumaric acid salt(6K)	0.009	0.04	6.40
1-(6-Bromo-3-pyridyl)-homopiperazine fumaric acid salt (8K)	0.001	0.01	0.62
1-(3-Pyridazinyl)-homopiperazine (1L)	0.03	0.25	1.60
1-(2-Quinoxaliny)-homopiperazine fumaric acid salt (2L)	0.26	2.00	27.00
1-(5-Methoxy-3-pyridyl)-3-methyl- imidazolidine fumaric acid salt (1M)	0.59	8.80	>30.0
1-(5-Methoxy-3-pyridyl)-3-methyl-1,3- diazacyclohexane fumaric acid salt (2M)	0.09	2.70	>30

Pharmaceutical Compositions

While it is possible that, for use in therapy, a compound of the invention may be administered as the raw chemical, it is preferable to present the active ingredient as a pharmaceutical formulation.

- 5 The invention thus further provides pharmaceutical formulations comprising a compound of the invention or a pharmaceutically acceptable salt or derivative thereof together with one or more pharmaceutically acceptable carriers therefor and, optionally, other therapeutic and/or prophylactic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

10

Pharmaceutical formulations include those suitable for oral, rectal, nasal, topical (including buccal and sub-lingual), vaginal or parenteral (including intramuscular, sub-cutaneous and intravenous) administration or in a form suitable for administration by inhalation or insufflation.

- 15 The compounds of the invention, together with a conventional adjuvant, carrier, or diluent, may thus be placed into the form of pharmaceutical compositions and unit dosages thereof, and in such form may be employed as solids, such as tablets or filled capsules, or liquids such as solutions, suspensions, emulsions, elixirs, or capsules filled with the same, all for oral use, in the form of suppositories for rectal administration; or in the form of sterile injectable solutions
- 20 for parenteral (including subcutaneous) use. Such pharmaceutical compositions and unit dosage forms thereof may comprise conventional ingredients in conventional proportions, with or without additional active compounds or principles, and such unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed. Formulations containing ten (10) milligrams of active ingredient
- 25 or, more broadly, 0.1 to one hundred (100) milligrams, per tablet, are accordingly suitable representative unit dosage forms.

The compounds of the present invention can be administered in a wide variety of oral and parenteral dosage forms. It will be obvious to those skilled in the art that the following dosage

30 forms may comprise, as the active component, either a compound of the invention or a pharmaceutically acceptable salt of a compound of the invention.

For preparing pharmaceutical compositions from the compounds of the present invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations

35 include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavouring agents,

solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

In powders, the carrier is a finely divided solid which is in a mixture with the finely divided
5 active component.

In tablets, the active component is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted in the shape and size desired.

- 10 The powders and tablets preferably contain from five or ten to about seventy percent of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as
15 carrier providing a capsule in which the active component, with or without carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid forms suitable for oral administration.
- 20 For preparing suppositories, a low melting wax, such as admixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by stirring. The molten homogenous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.
- 25 Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Liquid form preparations include solutions, suspensions, and emulsions, for example, water or
30 water-propylene glycol solutions. For example, parenteral injection liquid preparations can be formulated as solutions in aqueous polyethylene glycol solution.

The compounds according to the present invention may thus be formulated for parenteral administration (e.g. by injection, for example bolus injection or continuous infusion) and may
35 be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The compositions may take such forms as

suspensions, solutions, or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilisation from solution, for constitution with a suitable vehicle, e.g. sterile, pyrogen-free
5 water, before use.

Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavours, stabilizing and thickening agents, as desired.

- 10 Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, or other well known suspending agents.

Also included are solid form preparations which are intended to be converted, shortly before
15 use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavours, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

- 20 For topical administration to the epidermis the compounds according to the invention may be formulated as ointments, creams or lotions, or as a transdermal patch. Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilising agents, dispersing
25 agents, suspending agents, thickening agents, or colouring agents.

Formulations suitable for topical administration in the mouth include lozenges comprising active agent in a flavoured base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert base such as gelatin and glycerin or sucrose and
30 acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

Solutions or suspensions are applied directly to the nasal cavity by conventional means, for example with a dropper, pipette or spray. The formulations may be provided in single or multidose form. In the latter case of a dropper or pipette, this may be achieved by the patient
35 administering an appropriate, predetermined volume of the solution or suspension. In the case of a spray, this may be achieved for example by means of a metering atomising spray pump.

Administration to the respiratory tract may also be achieved by means of an aerosol formulation in which the active ingredient is provided in a pressurised pack with a suitable propellant such as a chlorofluorocarbon (CFC) for example dichlorodifluoromethane, 5 trichlorofluoromethane, or dichlorotetrafluoroethane, carbon dioxide, or other suitable gas. The aerosol may conveniently also contain a surfactant such as lecithin. The dose of drug may be controlled by provision of a metered valve.

Alternatively the active ingredients may be provided in the form of a dry powder, for example a 10 powder mix of the compound in a suitable powder base such as lactose, starch, starch derivatives such as hydroxypropylmethyl cellulose and polyvinylpyrrolidone (PVP). Conveniently the powder carrier will form a gel in the nasal cavity. The powder composition may be presented in unit dose form for example in capsules or cartridges of, e.g., gelatin, or blister packs from which the powder may be administered by means of an inhaler.

15 In formulations intended for administration to the respiratory tract, including intranasal formulations, the compound will generally have a small particle size for example of the order of 5 microns or less. Such a particle size may be obtained by means known in the art, for example by micronization.

20 When desired, formulations adapted to give sustained release of the active ingredient may be employed.

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the 25 preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

30 Tablets or capsules for oral administration and liquids for intravenous administration are preferred compositions.

Method of Treating

The compounds of the present invention are valuable nicotinic ACh receptor modulators and therefore useful for the treatment of a range of ailments involving cholinergic dysfunction as well as a range of disorders responsive to the activity of nicotinic ACh receptor modulators.

The compounds may be used in the treatment, prevention, prophylaxis or alleviation of a
5 disease, disorder or condition of the central or peripheral system as for example:
neurodegenerative disorders, cognitive or memory dysfunction, Alzheimer's disease,
Parkinson's disease, Huntington's disease, Amyotrophic Lateral Sclerosis, Gilles de la
Tourettes syndrome, attention deficit hyperactivity disorder, anxiety, depression, mania, manic
depression, schizophrenia, obsessive compulsive disorders, eating disorders like anorexia
10 nervosa, bulimia and obesity, narcolepsy, nociception, memory loss, memory dysfunction,
AIDS-dementia, senile dementia, peripheral neuropathy, learning deficit, cognition deficit,
attention deficit, autism, dyslexia, tardive dyskinesia, hyperkinesia, epilepsy, bulimia, post-
traumatic syndrome, social phobia, chronic fatigue syndrome, disorders of sleep,
pseudodementia, Ganser's syndrome, premenstrual syndrome, late luteal phase syndrome,
15 chronic fatigue syndrome, premature ejaculation, erectile difficulty, mutism and
trichotillomania.

The compounds of this invention may also be used in the treatment of inflammatory conditions
as for example: inflammatory skin conditions like acne and rosacea, Chron's disease,
20 inflammatory bowel disease, ulcerative colitis, irritable colon, diarrhoea.

Also the compounds of the invention may be used in the treatment of diseases associated with
smooth muscle contractions as for example: convulsive disorders, angina pectoris, premature
labor, convulsions, diarrhoea, asthma, epilepsy, tardive dyskinesia, hyperkinesia.
25

The compounds of this invention may also be used in the treatment of pain as for example
chronic, acute and recurrent pain, postoperative pain, migraine pain or phantom limb pain;
The compounds of the present invention may also be used for the assistance in cessation of
abuse of chemical substances as for example smoking cessation as well as cessation of use
30 of other nicotine containing products, cessation of use of opioids like heroin, cocaine and
morphine and cessation of use of benzodiazepines or alcohol. In the context of the present
invention "treatment" means as well treatment as prevention, prophylaxis and alleviation of
withdrawal symptoms and abstinence as well as treatment resulting in a voluntary diminished
intake of the addictive substance.

Suitable dosage range are 0.1-500 milligrams daily, and especially 10-70 milligrams daily, administered once or twice a day, dependent as usual upon the exact mode of administration, form in which administered, the indication toward which the administration is directed, the subject involved and the body weight of the subject involved, and further the preference and
5 experience of the physician or veterinarian in charge.

I.p. means intraperitoneally, which is a well known route of administration.

P.o. means peroral, which is a well known route of administration.

10 The invention then comprises the following alone or in combination:

The use as above wherein the disease to be treated is pain, a disease in the central or peripheral system, a disease caused by smooth muscle contraction, neurodegeneration, inflammation, chemical substance abuse or withdrawal symptoms caused by the cessation of
15 intake of the chemical substance, such as assistance in the cessation of smoking;

The use as above wherein a disease in the central or peripheral system is Alzheimer's disease, Parkinson's disease, memory dysfunction or attention deficit hyperactivity disorder.

20 The method as above wherein pain, a disease in the central or peripheral system, a disease caused by smooth muscle contraction, neurodegeneration, inflammation, chemical substance abuse or withdrawal symptoms caused by the cessation of intake of chemical substances, such as smoking cessation, is treated.

The method as above wherein chemical substance abuse or withdrawal symptoms caused by
25 the cessation of intake of the chemical substance, said chemical substance abuse being smoking or use of other nicotine containing products and withdrawal symptoms caused by cessation of use of nicotine containing products, is treated;

A method as above wherein a disease in the central or peripheral system, said disease being Alzheimer's disease, Parkinson's disease, memory dysfunction or attention deficit hyperactivity
30 disorder, is treated.

The following examples will illustrate the invention further, however, they are not to be construed as limiting.

Examples

General: All reactions involving air sensitive reagents or intermediates were performed under nitrogen and in anhydrous solvents. Magnesium sulfate is used as drying agent in the workup-procedures and solvents were evaporated under reduced pressure.

5 Method A

4-Methyl-1-(3-pyridyl)-piperazine fumaric acid salt (1A)

A solution of 1-(3-pyridyl)-piperazine (0.35 g, 2.1 mmol), formic acid (1.0 g, 21.7 mmol), formaldehyde (0.64 g, 37 %) and water (2 ml) was stirred at reflux for 15 h. The mixture was evaporated and sodium hydroxide (30 ml, 1 M) was added and the product was extracted
10 three times with ethyl acetate (15 ml). The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Yield 0.21 g, 34 %. Mp 144.5 - 145.9 °C.

4-Methyl-1-(3-quinoliny)-piperazine (2A)

15 Was prepared according to method A. Isolated as free base. Mp 116.5 - 117.0 °C. This compound is previously known in the literature (B. Schonen and F. Zymalkowski, Arch Pharm. (Weinheim) 314, 464 - 470 (1981)).

4-Methyl-1-(5-methoxy-3-pyridyl)-piperazine (3A)

20 Was prepared according to method A. Isolated as free base. Mp 67.5 - 68.0 °C.

3,5-Bis-[4,4'-methyl-1,1'-piperazinyl]pyridine (4A)

Was prepared according to method A. Mp 132 - 133 °C.

25 1-(5-Chloro-3-pyridyl)-4-Methyl-piperazine fumaric acid salt (5A)

Was prepared according to method A. Mp 162 - 163 °C.

4-Methyl-1-(5-phenyl-3-pyridyl)-piperazine fumaric acid salt (6A)

Was prepared according to method A. Mp 179 - 180 °C.

30

1-(5-Ethoxy-3-pyridyl)-4-methyl-piperazine fumaric acid salt (7A)

Was prepared according to method A. Mp 162.2 - 163.7 °C.

1-(5-Butoxy-3-pyridyl)-4-methyl-piperazine fumaric acid salt (8A)

35 Was prepared according to method A. Mp 136.9 - 139.2 °C.

trans-4-Methyl-1-[5-(Propyl-1-en-oxy)-3-pyridyl]-piperazine fumaric acid salt (9A)

Was prepared according to method A. Mp 145.1 - 145.7 °C.

5 1-(5-Ethenyloxy-3-pyridyl)-4-methyl-piperazine fumaric acid salt (10A)

Was prepared according to method A. Mp 136.4 - 138.2 °C.

1-(5-Methoxy-3-pyridyl)-4-methyl-1,5-diazacyclooctane fumaric acid salt (11A)

Was prepared according to method A. Mp 137 - 139 °C.

10

1-(6-Chloro-3-pyridazinyl)-4-methyl-1,5-diazacyclooctane fumaric acid salt (12A)

Was prepared according to method A. Mp 166 - 168 °C.

Method B**15 1-(3-Pyridyl)-piperazine fumaric acid salt (1B)**

A solution of 1-(3-pyridyl)-4-*tert*-butoxycarbonylpiperazine (1.3 g, 4.9 mmol), trifluoroacetic acid (11.3 g, 99 mmol) and dichloromethane (50 ml) was stirred for 15 h. The mixture was evaporated. Sodium hydroxide (4 M) was added. The product was extracted three times with dichloromethane (50 ml) and isolated as an oil. Yield 0.72 g, 90 %. The corresponding salt
20 was obtained by addition of a diethyl ether and methanol mixture (9:1), saturated with fumaric acid. Mp 161.7 - 164.8 °C.

1-(3-Quinoliny)-piperazine (2B)

Was prepared according to method B. The reactant was obtained according to method C from
25 3-bromoquinoline. Mp 87.7 - 88.5 °C.

1-(5-Methoxy-3-pyridyl)-piperazine fumaric acid salt (3B)

Was prepared according to method B. The reactant was obtained according to method C from 3-bromo-5-methoxypyridine. Mp 168.5 - 170.5 °C.

30

1-(5-Chloro-3-pyridyl)-piperazine fumaric acid salt (4B)

Was prepared according to method B. The reactant was obtained according to method C from 3,5-dichloropyridine. Mp 195 - 196 °C.

35 1-(5-Phenyl-3-pyridyl)-piperazine fumaric acid salt (5B)

Was prepared according to method B. The reactant was obtained according to method C from 3-bromo-5-phenylpyridine. Mp 167.5 - 168.5 °C.

1-(5-Methoxy-3-pyridyl)-1,5-diazacyclooctane fumaric acid salt (6B)

- 5 Was prepared according to method B. The reactants were obtained according to method C from 3-bromo-5-methoxypyridine and 1-*tert*-butoxycarbonyl-(1,5-diazacyclooctane). Mp 158 - 160 °C.

1-(6-Chloro-3-pyridazinyl)-1,5-diazacyclooctane fumaric acid salt (7B)

- 10 A mixture of 1,5-diazacyclooctane (2.07 g, 18.1 mmol), 3,6-dichloropyridazine (2.70 g, 18.1 mmol) and toluene (50 ml) was stirred at reflux overnight. Aqueous sodium hydroxide (50 ml, 1 M) was added followed by extraction seven times with ethyl acetate (50 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound. Yield 1.72 g, 42 %. The corresponding salt was obtained by addition of a diethyl
15 ether and methanol mixture (9 : 1) saturated with fumaric acid. Mp 176 - 178 °C.

1-(6-Chloro-3-pyridazinyl)-1,4-diazacyclooctane fumaric acid salt (8B)

- A mixture of 1,4-diazacyclooctane (2.07 g, 18.1 mmol), (this starting material was prepared according to J. Hernandez-Mora and Nadia Cordero-Antunano, Carib. J. Sci., 14, 77, 1974)
20 3,6-dichloropyridazine (2.70, 18.1 mmol) and toluene (50 ml) was stirred at reflux overnight. Aqueous sodium hydroxide (50 ml, 1 M) was added followed by extraction seven times with ethyl acetate (40 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound. Yield 1.2 g, 29 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9 : 1) saturated with fumaric
25 acid. Mp 177 - 179 °C.

Method C

1-(3-Pyridyl)-4- *tert*-butoxycarbonyl-piperazine

- A mixture of 3-bromopyridine (7.8 g, 49.4 mmol), 1-*tert*-butoxycarbonylpiperazine (9.2 g, 49.4
30 mmol), tetrakis(triphenylphosphine)palladium(0) (286 mg, 0.247 mmol), potassium *tert*-butoxide (11.1 g, 98.8 mmol) and anhydrous toluene (100 ml) was stirred at 80 °C for 0.5 h. Water (100 ml) was added and the mixture was extracted three times with ethyl acetate (75 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound as an oil. Yield 1.34 g, 10 %.

Method D**1-(5-Ethoxy-3-pyridyl)piperazine fumaric acid salt (1D)**

A mixture of 3-chloro-5-ethoxypyridine (6.5 g, 45.8 mmol), piperazine (19.7 g, 229 mmol), potassium *tert*-butoxide (11.2 g, 91.6 mmol) and 1,2-dimethoxyethane (150 ml) was stirred at
5 reflux for 1 h. Aqueous sodium hydroxide (1 M, 100 ml) was added and the mixture was
extracted two times with ethyl acetate (150 ml). Chromatography on silica gel with
dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound. Yield 4.6 g,
48 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture
(9:1) saturated with fumaric acid. Mp 160.0 - 161.2 °C.

10

1-(5-Butoxy-3-pyridyl)piperazine fumaric acid salt (2D)

Was prepared according to method D. Mp 149.4 - 151.7 °C.

***trans*-1-[5-(Propyl-1-en-oxy)-3-pyridyl]-piperazine fumaric acid salt (3D)**

15 Was prepared according to method D, from 3-chloro-5-(propyl-2-en-oxy)-pyridine involving
isomerization of the double bond. Mp 145.1 - 145.7 °C.

1-(5-Ethenyloxy-3-pyridyl)piperazine fumaric acid salt (4D)

Was prepared according to method D. Mp 136.4 - 138.2 °C.

20

1-(5-Ethoxy-3-pyridyl)-1,5-diazacyclooctane fumaric acid salt (5D)

Was prepared according to method D from 1,5-diazacyclooctane (prepared according to J.
Hernandez-Mora "Derivatives of 1,5-diazacyclooctane " Ph.D. Dissertation, University of
Michigan (1959)) at room temperature over night. Mp 162.5 - 164.5 °C.

25

1-(5-Propyloxy-3-pyridinyl)-1,5-diazacyclooctane fumaric acid salt (6D)

Was prepared according to method D from 1,5-diazacyclooctane at 70 °C over night in the
presence of tetrakis(triphenylphosphine)palladium(0) (3 %). Mp 150 - 152 °C.

30 ***trans*-1-[5-(Propyl-1-en-oxy)-3-pyridyl]-1,5-diazacyclooctane fumaric acid salt (7D)**

Was prepared according to method D, from 3-chloro-5-(propyl-2-en-oxy)-pyridine involving
isomerization of the double bond. Mp 135 - 137 °C.

1-(5-Ethenyloxy-3-pyridyl) 1,5-diazacyclooctane fumaric acid salt (8D)

35 Was prepared according to method D. Mp 142 - 144 °C.

1-(5-Ethoxy-3-pyridyl)-4-ethyl-piperazine fumaric acid salt (9D)

A mixture of 1-(5-ethoxy-3-pyridyl)-piperazine (1.4 g, 6.8 mmol), triethylamine (0.69 g, 6.8 mmol), bromoethane (0.88 g, 8.1 mmol) and dimethylformamide (25 ml). Aqueous sodium hydroxide (1 M, 50 ml) was added and the mixture was extracted two times with ethyl acetate (50 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound. Yield 0.75 g, 47 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1), saturated with fumaric acid. Mp 144.8 - 145.9 °C.

10

General procedure for 3-bromo and 3-chloro-5-arylpyridines:**3-Bromo-5-phenylpyridine**

A mixture of 3,5-dibromopyridine (10.0 g, 42.2 mmol), phenylboronic acid (4.6 g, 38.0 mmol), tetrakis(triphenylphosphine)palladium(0) (1.45 g, 1.25 mmol), potassium carbonate (17.5 g, 127 mmol), water (63 ml) and 1,2-dimethoxyethane (126 ml) was stirred at reflux overnight. Aqueous sodium hydroxide (1 M, 60 ml) was added followed by extraction twice with diethyl ether (100 ml). Chromatography on silica gel with dichloromethane as solvent gave the title compound. Yield 6.1 g, 68 %, Mp 42 - 44 °C.

20

3-Bromo-6-thioethoxypyridine

A mixture of sodium thioethoxide (7.81 g, 92.9 mmol), 2,5-dibromopyridine (20.0 g, 84.4 mmol) and dimethyl sulfoxide (100 ml). The mixture was stirred at 20 °C overnight. Sodium hydroxide (300 ml, 1 M) was added and the mixture was extracted twice with diethyl ether (200 ml). Chromatography on silica gel with dichloromethane : petroleum ether, 1:2 as eluent, gave the title compound as an oil. Yield 16.8 g, 85 %.

Method E**4-Methyl-1-(3-pyridyl)-homopiperazine (1E)**

A solution of 1-(3-pyridyl)-homopiperazine (0.42 g, 2.4 mmol), formic acid (3.3 g, 71.7 mmol), formaldehyde (2.1 g, 37 %) and water (10 ml) was stirred at reflux for 15 h. The mixture was evaporated and sodium hydroxide (15 ml, 4 M) was added and the product was extracted two times with ethyl acetate (15 ml). The product was obtained as an oil. Yield 0.46 g, 100 %.

35 4-Methyl-1-(3-quinolinyl)-homopiperazine fumaric acid salt (2E)

Was prepared according to method E. Mp 170 - 171 °C.

1-(5-Methoxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (3E)

Was prepared according to method E. Mp 145 - 147 °C.

5

1-(5-Ethoxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (4E)

Was prepared according to method E. Mp 150 - 152 °C.

4-Methyl-1-(5-phenyl-3-pyridyl)-homopiperazine fumaric acid salt (5E)

10 Was prepared according to method E. Mp 161 - 162 °C.

1-(5-Butoxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (6E)

Was prepared according to method E. Mp 127 - 129 °C.

15 **1-(5-Methoxyethoxy-3-pyridyl)-4-methyl-homopiperazine (7E)**

Was prepared according to method E. Isolated as an oil.

4-Methyl-1-[5-(2-methyl-propyloxy)-3-pyridyl]-homopiperazine fumaric acid salt (8E)

Was prepared according to method E. Mp 145.7 - 146.9 °C.

20

1-(5-Cyclopropylmethoxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (9E)

Was prepared according to method E. Mp 160.4 - 161.9 °C.

4-Methyl-1-(5-propyloxy-3-pyridyl)-homopiperazine fumaric acid salt (10E)

25 Was prepared according to method E. Mp 148.8 - 153.5 °C.

1-(5-Hexyloxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (11E)

Was prepared according to method E. Mp 128.7 - 130.8 °C.

30 **4-Methyl-1-[5-(3-methyl-butoxy)-3-pyridyl]-homopiperazine fumaric acid salt (12E)**

Was prepared according to method E. Mp 130.4 - 131.9 °C.

4-Methyl-1-(6-thioethoxy-3-pyridyl)-homopiperazine fumaric acid salt (13E)

Was prepared according to method E. Mp 119 - 121 °C.

1-(5-Cyclohexylmethoxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (14E)

Was prepared according to method E. Mp 160.4 - 162.0 °C.

4-Methyl-1-(5-pentyloxy-3-pyridyl)-homopiperazine fumaric acid salt (15E)

5 Was prepared according to method E. Mp 129.0 - 130.8 °C.

1-(5-Heptyloxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (16E)

Was prepared according to method E. Mp 120.2 - 121.8 °C.

10 ***trans*-4-Methyl-1-(5-propyl-1-en-oxy-3-pyridyl)-homopiperazine fumaric acid salt (17E)**

Was prepared according to method E. Mp 126 - 128 °C.

4-Methyl-1-(5-thiobenzyl-3-pyridyl)-homopiperazine fumaric acid salt (18E)

Was prepared according to method E. Mp 131 - 133 °C.

15

4-Methyl-1-[5-(3-pyridyl)-3-pyridyl]-homopiperazine fumaric acid salt (19E)

Was prepared according to method E. Mp 165.5 - 167.5 °C.

4-Methyl-1-(3-nitrophenyl)-homopiperazine (20E)

20 Was prepared according to method E. Mp 163.1 - 164.4 °C.

4-Methyl-1-(6-chloro-3-pyridazinyl)-homopiperazine fumaric acid salt (21E)

Was prepared according to method E. Mp 171 - 172 °C.

25 **4-Methyl-1-(6-phenyl-3-pyridazinyl)-homopiperazine fumaric acid salt (22E)**

Was prepared according to method E. Mp 185 - 186 °C.

4-Methyl-1-(3-pyridazinyl)-homopiperazine fumaric acid salt (23E)

Was prepared according to method E. Mp 137.8 - 139.3 °C.

30

4-Methyl-1-(6-methyl-3-pyridazinyl)-homopiperazine fumaric acid salt (24E)

Was prepared according to method E. Mp 152 - 153 °C.

1-(5-Cyclopentyloxy-3-pyridyl)-4-methyl-homopiperazine fumaric acid salt (25E)

35 Was prepared according to method E. Mp 123 - 125 °C.

4-Benzyl-1-(3-Pyridyl)-homopiperazine fumaric acid salt (26E)

1-(3-Pyridyl)-homopiperazine (0.54 g, 3.0 mmol), potassium carbonate (0.42 g, 3.0 mmol), benzylbromide (0.56 g, 3.3 mmol) in dimethylformamide (40 ml) was stirred at 80 °C for one
5 hour. Water (100 ml) was added and the mixture was extracted twice with ethyl acetate (25 ml). Yield 0.39 g, 49 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1), saturated with fumaric acid. Mp 148.4 - 149.0 °C.

4-Ethyl-1-(3-pyridyl)-homopiperazine fumaric acid salt (27E)

10 Was prepared according to 4-benzyl-1-(3-pyridyl)-homopiperazine from 1-(3-pyridyl)homopiperazine. Mp 145.3 - 147.5 °C.

Method F**1-(3-Pyridyl)-homopiperazine fumaric acid salt (1F)**

15 A solution of 1-(3-pyridyl)-4-*tert*-butoxycarbonylhomopiperazine (0.91 g, 3.3 mmol), trifluoroacetic acid (7.5 g, 66 mmol) and dichloromethane (30 ml) was stirred for 15 h. The mixture was evaporated. Sodium hydroxide (30 ml, 4 M) was added. The product was extracted two times with dichloromethane (30 ml) and isolated as an oil. Yield 0.50 g, 85 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1)
20 saturated with fumaric acid. Mp 172.1 - 172.9 °C.

1-(3-Quinoliny)-homopiperazine fumaric acid salt (2F)

Was prepared according to method F. Mp 181 - 182 °C.

1-(6-Methoxy-3-pyridyl)-homopiperazine fumaric acid salt (3F)

Was prepared according to method F. Mp 127 - 128 °C.

1-(2-Chloro-5-methoxy-3-pyridyl)-homopiperazine fumaric acid salt (4F)

Was prepared according to method F. Mp 117 - 118 °C.

30

1-(6-Chloro-5-methoxy-3-pyridyl)-homopiperazine fumaric acid salt (5F)

Was prepared according to method F. Mp 196 - 197 °C.

1-(6-Bromo-5-ethoxy-3-pyridyl)-homopiperazine fumaric acid salt (6F)

35 Was prepared according to method F. Mp 181.7 - 183.2 °C.

1-[6-(N-Pyrrolidinyl)-3-pyridyl]-homopiperazine fumaric acid salt (7F)

Was prepared according to method F. Mp 148.6 - 150.5 °C.

5 1-(6-Phenyl-3-pyridyl)-4-homopiperazine fumaric acid salt (8F)

Was prepared according to method F. Mp 180 - 182 °C.

1-[5-(3-Nitrophenyl)-3-pyridyl]-4-homopiperazine fumaric acid salt (9F)

Was prepared according to method F. Mp 186 - 188 °C.

10

Method G**1-(3-Pyridyl)-4-*tert*-butoxycarbonylhomopiperazine**

A mixture of 3-bromopyridine (3.95 g, 25.0 mmol), 1-*tert*-butoxycarbonylhomopiperazine (5.0 g, 25.0 mmol), tetrakis(triphenylphosphine)palladium(0) (145 mg, 0.125 mmol), potassium *tert*-
15 butoxide (6.1 g, 50.0 mmol) and anhydrous toluene (75 ml) was stirred at 80 °C for 4 h. Water (100 ml) was added and the mixture was extracted three times with ethyl acetate (50 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound as an oil. Yield 0.92 g, 13 %.

20 **1-(6-Methoxy-3-pyridyl)-4-*tert*-butoxycarbonylhomopiperazine** Was prepared according to Method G. Isolated as an oil.

1-(3-Quinoliny)-4-*tert*-butoxycarbonylhomopiperazine

Was prepared according to Method G. Isolated as an oil.

25

Method H**1-(5-Methoxy-3-pyridyl)-homopiperazine fumaric acid salt (1H)**

A mixture of 3-bromo-5-methoxypyridine (5.6 g, 30.0 mmol), homopiperazine (15.0 g, 150 mmol), tetrakis(triphenylphosphine)palladium(0) (173 mg, 0.15 mmol), potassium-*tert*-butoxide
30 (6.7 g, 60 mmol) and anhydrous toluene (150 ml) was stirred at 80 °C for 4 h. Water (100 ml) was added and the mixture was extracted seven times with ethyl acetate (150 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound. Yield 3.5 g, 56 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 161 - 162 °C.

35

1-(5-Phenyl-3-pyridyl)-homopiperazine fumaric acid salt (2H)

Was prepared according to method H. Mp 185 - 186 °C.

1-(5-Ethoxy-3-pyridyl)-homopiperazine fumaric acid salt (3H)

5 Was prepared according to method H. Mp 157.5 - 159 °C.

1-(5-Butoxy-3-pyridyl)-homopiperazine fumaric acid salt (4H)

Was prepared according to method H. Mp 150 - 151 °C.

10 **1-(5-Methoxyethoxy-3-pyridyl)-homopiperazine fumaric acid salt (5H)**

Was prepared according to method H. Mp 126 - 127 °C.

1-[5-(2-Methyl-propoxy)-3-pyridyl]-homopiperazine fumaric acid salt (6H)

Was prepared according to method H. Mp 121.9 - 123.3 °C.

15

1-[5-(2-Hydroxy-ethoxy)-3-pyridyl]-homopiperazine (7H)

Was prepared according to method H. Was isolated as an oil.

1-[5-(3-Methyl-butoxy)-3-pyridyl]-homopiperazine fumaric acid salt (8H)

20 Was prepared according to method H. Mp 139.9 - 142.0 °C.

1-(5-Cyclopropylmethoxy-3-pyridyl)-homopiperazine fumaric acid salt (9H)

Was prepared according to method H. Mp 154 - 156 °C.

25 **1-(5-Propyloxy-3-pyridyl)-homopiperazine fumaric acid salt (10H)**

Was prepared according to method H. Mp 156.2 - 157.8 °C.

1-(5-Hexyloxy-3-pyridyl)-homopiperazine fumaric acid salt (11H)

Was prepared according to method H. Mp 149.5 - 151.8 °C.

30

1-(5-Cyclohexylmethoxy-3-pyridyl)-homopiperazine fumaric acid salt (12H)

Was prepared according to method H. Mp 163.3 - 164.5 °C.

1-(6-Thioethoxy-3-pyridyl)-homopiperazine fumaric acid salt (13H)

35 Was prepared according to method H. Mp 115 - 119 °C.

1-[5-(2-Ethoxy-ethoxy)-3-pyridyl]-homopiperazine fumaric acid salt (14H)

Was prepared according to method H. Mp 139.3 - 140.4 °C.

5 1-(5-Pentyloxy-3-pyridyl)-homopiperazine fumaric acid salt (15H)

Was prepared according to method H. Mp 155.5 - 156.7 °C.

1-(5-Heptyloxy-3-pyridyl)-homopiperazine fumaric acid salt (16H)

Was prepared according to method H. Mp 132.8 - 136.6 °C.

10

***trans*-1-[5-(Propyl-1-en-oxy)-3-pyridyl]-homopiperazine fumaric acid salt (17H)**

Was prepared according to method H, from 3-chloro-5-(propyl-2-en-oxy)-pyridine involving isomerization of the double bond. Mp 124 -126 °C.

15 1-(5-Thiobenzyl-3-pyridyl)-homopiperazine fumaric acid salt (18H)

Was prepared according to method H. Mp 148 - 150 °C.

1-(5-Carboxylamido-3-pyridyl)-homopiperazine fumaric acid salt (19H)

Was prepared according to method H. Mp 149 - 151 °C.

20

1-(5-Thiophenyl-3-pyridyl)-homopiperazine fumaric acid salt (20H)

Was prepared according to method H. Mp 177 - 179 °C.

Method I**25 1-[(5-Methoxy-methoxy)-3-pyridyl]-homopiperazine fumaric acid salt (1I)**

A mixture of 3-chloro-5-methoxymethoxypyridine (10.0 g, 57.6 mmol), homopiperazine (28.8 g, 288 mmol), 1.3-bis(diphenylphosphino)propanepalladiumdichloride (170 mg, 0.29 mmol), potassium *tert*-butoxide (12.9 g, 115 mmol) and 1,2-dimethoxyethane (100 ml) was stirred at reflux for 3 h. Sodium hydroxide (1 M, 100 ml) was added and the mixture was extracted two
30 times with ethyl acetate (150 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound as free base. Yield 9.7 g, 71 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 129.5 - 131 °C.

35 1-[5-(3-Pyridyl)-3-pyridyl]-homopiperazine fumaric acid salt (2I)

Was prepared according to method I. Mp 160 - 162 °C.

Method J

1-(5-(1-Pyrrolyl)-3-pyridyl)-homopiperazine fumaric acid salt (1J)

- 5 A mixture of 3-chloro-5-(1-pyrrolyl)-pyridine (6.3 g, 35.3 mmol), homopiperazine (7.06 g, 70.5 mmol), potassium *tert*-butoxide (7.91 g, 70.5 mmol) and 1,2-dimethoxyethane (100 ml) was stirred at reflux for 3 h. Sodium hydroxide (1 M, 120 ml) was added and the mixture was extracted three times with ethyl acetate (100 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound. Yield 3.45
10 g, 40 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 174 - 175 °C.

1-(5-(1-Indolyl)-3-pyridyl)-homopiperazine fumaric acid salt (2J)

Was prepared according to method J from 3-chloro-5-(1-indolyl)-pyridine. Mp 193 - 195 °C.

15

1-(5.6-Dimethoxy-3-pyridyl)-homopiperazine fumaric acid salt (3J)

Was prepared according to method J from 3-chloro-5.6-dimethoxypyridine. Mp 150 - 152 °C.

1-(5-Ethenyloxy-3-pyridyl)homopiperazine fumaric acid salt (4J)

- 20 Was prepared according to method J from 3-Chloro-5-ethenyloxypyridine. Mp 143 -144 °C.

1-(5-Cyclopentyloxy-3-pyridyl)homopiperazine fumaric acid salt (5J)

Was prepared according to method J. Mp 148 -150 °C.

- 25 **1-[5-(Ethoxy-d5)-3-pyridyl]-homopiperazine fumaric acid salt (6J)**

Was prepared according to method J, using 3-chloro-5-(penta-deuterium-ethoxy)-pyridine as starting material. Mp 163 - 165 °C.

1-(3-Chloro-5-pyridyl)homopiperazine fumaric acid salt (7J)

- 30 Was prepared according to method J, using room temperature as reaction temperature. Mp 144.4 -146.6 °C.

1-(3-Bromo-5-pyridyl)homopiperazine fumaric acid salt (8J)

Was prepared according to method J, using room temperature as reaction temperature. Mp

- 35 180.7 -185.4 °C.

1-(4-Isoquinolinyl)-homopiperazine fumaric acid salt (9J)

A mixture of 4-bromoisoquinoline (0.80 g, 3.85 mmol) and homopiperazine (3.85 g, 38.5 mmol) was stirred at 170 °C overnight. Aqueous sodium hydroxide (20 ml, 1M) was added.

- 5 The mixture was extracted three times with ethyl acetate (30 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound. Yield 0.40 g, 46 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 160 - 162 °C.

10 **3-Chloro-5-(1-pyrrolyl)-pyridine**

A mixture of 3,5-dichloropyridine (10.0 g, 67.6 mmol), pyrrole (5.50 g, 81.1 mmol), sodium hydride 60 % (3.52 g, 87.9 mmol) and dimethyl sulfoxide (50 ml) was stirred at 70 °C for 2 h. Aqueous sodium hydroxide (200 ml, 1 M) was added and the mixture was extracted three times with diethyl ether (100 ml). The mixture was evaporated and, purified by

- 15 chromatography on silica gel with dichloromethane and ethanol (4 %) as solvent. Yield 6.3 g, 52 %. Mp 70.5 - 72.0 °C.

3-Chloro-5-(1-indolyl)-pyridine

Was prepared according to 3-Chloro-5-(1-pyrrolyl)-pyridine. Yield 5.9 g, 38 %. Mp 56 - 57 °C.

20

3-Chloro-5-ethenyloxy-pyridine

Thionyl chloride (58.6 g, 492.6 mmol) was added to a mixture of 1-[3-chloro-5-(2-hydroxyethoxy)]pyridine (14.5 g, 82.1 mmol) and tetrahydrofuran (100). The mixture was stirred for 1 h at 50 °C. The mixture was evaporated. Potassium hydroxide (9.0 g, 164 mmol)

- 25 and *tert*-butanol (100 ml) was added and the mixture was stirred for 3 days at 100 °C. The solvent was evaporated. Water (150 ml) was added and extracted twice with diethyl ether (100 ml). Yield 6.77 g, 53 %.

1-(5-Hydroxy-3-pyridyl)-homopiperazine hydrochloric acid salt (10J)

- 30 1-(5-Methoxymethoxy-3-pyridyl)-homopiperazine (8.5 g, 35.9 mmol) was stirred in hydrochloric acid (4 M, 100 ml) at room temperature for 1 h. The excess of hydrochloric acid was evaporated. A crystalline compound was obtained by triturating with a mixture of 5% methanol and ether. Yield 9.56 g, 100 %. Mp 290 - 300 °C.

- 35 **1-(5-Trifluoromethanesulfonyl-oxy-3-pyridyl)-homopiperazine fumaric acid salt (11J)**

To a mixture of 1-(5-trifluoromethanesulfonyl-oxy-3-pyridyl)-4-*tert*-butoxycarbonyl-homopiperazine (0.82 g, 1.9 mmol) and dichloromethane (10 ml) was added trifluoro acetic acid (2.18 g, 19.2 mmol) at room temperature. The mixture was stirred at room temperature for 3 h. Aqueous sodium hydroxide (30 ml) was added and the mixture was extracted twice
5 with dichloromethane (30 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 154 - 156 °C. Yield 0.29 g, 35 %.

10 1-(5-Trifluoromethanesulfonyl-oxy-3-pyridyl)-4-*tert*-butoxycarbonyl-homopiperazine

To a mixture of 1-(5-hydroxy-3-pyridyl)-4-*tert*-butoxycarbonyl-homopiperazine (4.0 g, 13.6 mmol), pyridine (3.23 g, 40.8 mmol) and dichloromethane (40 ml) was added: trifluoromethane sulfonic anhydride (3.85 g, 13.6 mmol) at 0 °C. The mixture was allowed to react at room temperature overnight. The organic phase was washed twice with aqueous sodium hydroxide (1
15 M, 30 ml). Chromatography on silica gel with ethyl acetate - toluene (2:1). Yield 2.26 g, 56 %.

1-(5-Hydroxy-3-pyridyl)-4-*tert*-butoxycarbonyl-homopiperazine

A mixture of 1-(5-hydroxy-3-pyridyl)-homopiperazine hydrochloric acid salt (15.5 g, 58.1 mmol) *tert*-butoxycarbonyl anhydride (12.7 g, 58.1 mmol), an aqueous solution of sodium hydrogen
20 carbonate (1 M, 290 mmol) and dichloromethane (290 ml) was stirred overnight. The organic phase was separated and purified with chromatography on silica gel, using 6 % ethanol and dichloromethane as eluent. The product was isolated as an oil. Yield 8.17 g, 48 %.

1-(5-Ethynyl-3-pyridyl)-homopiperazine fumaric acid salt (12J)

25 A mixture of 1-(5-ethynyl-3-pyridyl)-4-*tert*-butoxycarbonyl-homopiperazine (0.13 g, 0.43 mmol), trifluoro acetic acid (0.98 g, 8.6 mmol) and dichloromethane (10 ml) was stirred for 5 h. Aqueous sodium hydroxide (1 M, 15 ml) was added the organic phase was separated and the aqueous phase was extracted twice with dichloromethane (15 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound.
30 The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Yield 30 mg, 22 %. Mp 172.5 - 174.0 °C.

1-(5-Ethynyl-3-pyridyl)-4-*tert*-butoxycarbonyl-homopiperazine

A mixture of 1-[5-(3-methyl-3-hydroxy-butyn-1-yl)-3-pyridyl]-4-*tert*-butoxycarbonyl-
35 homopiperazine (0.40 g, 1.1 mmol), sodium hydride 60 % (4.5 mg, 0.11 mmol) and toluene

(10 ml) was stirred at 110 °C for 3 h. The crude mixture was purified by chromatography on silica gel with ethyl acetate : toluene, (3 : 1) gave the title compound. Yield 0.13 g, 39 %.

1-[5-(3-Methyl-3-hydroxy-butyn-1-yl)-3-pyridyl]-4-*tert*-butoxycarbonyl-homopiperazine

- 5 A mixture of 1-(5-trifluoromethanesulfonyl-oxy-3-pyridyl)-4-*tert*-butoxycarbonyl-homopiperazine (8.15 g, 1.9 mmol), potassium carbonate (0.66 g, 4.8 mmol), copper (I)iodide (37 mg, 0.19 mmol), palladium on carbon (5 %, 10 mg), triphenylphosphine (50 mg, 0.19 mmol), lithium chloride (81 mg, 1.9 mmol) and 1,2-dimethoxyethane (15 ml) was stirred at room temperature for 30 min. 2-Methyl-3-butyn-2-ol (1.62 g, 3.8 mmol) solved in 1,2-
- 10 dimethoxyethane (30 ml) was added to the mixture which was stirred at reflux overnight. The crude mixture was filtered through celite and hydrochloric acid (20 ml, 2 M) and toluene (30 ml) was added. The organic phase was discarded and the mixture was made alkaline with aqueous sodium hydroxide, followed by extraction with ethyl acetate (30 ml). The mixture was purified by chromatography on silica gel with ethyl acetate : toluene, (3:1) and gave the title
- 15 compound. Yield 0.40 g, 58 %.

1-(2-Chloro-5-methoxy-3-pyridyl)-4-*tert*-butoxycarbonylhomopiperazine and 1-(6-Chloro-5-methoxy-3-pyridyl)-4-*tert*-butoxycarbonylhomopiperazine

- An aqueous solution of sodium hypochlorite (16.3 ml, 8.14 mmol) was added to mixture of 1-
- 20 (5-methoxy-3-pyridyl)-4-*tert*-butoxycarbonylhomopiperazine (2.5 g, 8.14 mmol) and dimethylformamide (185 ml) at room temperature and stirred for 0.5 h at room temperature. Water was added (300 ml) and the mixture was extracted twice with diethyl ether (200 ml). The mixture was separated by chromatography on silica gel with ethyl acetate : toluene (2:1). The title compounds eluated in the order indicated above in in 2.0 g and 0.5 g respectively,
- 25 total yield 90 %.

1-[5-(3-Nitrophenyl)-3-pyridyl]-4-*tert*-butoxycarbonylhomopiperazine (13J)

- A mixture of (5-trifluoromethanesulfonyl-oxy-3-pyridyl)-4-*tert*-butoxycarbonyl-homopiperazine (3.0 g, 7.1 mmol), 1,3-propandiol (2.68 g, 35.3 mmol), lithium chloride (0.90 g, 21.2 mmol),
- 30 potassium carbonate (10.6 ml, 2 M), tetrakis(triphenylphosphine)palladium(0) (244 mg, 0.21 mmol), 3-nitrophenyl boronic acid (1.77 g, 10.6 mmol) and 1,2-dimethoxyethane was stirred at reflux for 2 h. Aqueous sodium hydroxide was added. The mixture was extracted twice with ethyl acetate (40 ml). The mixture was separated by chromatography on silica gel with petroleum : ethyl acetate, (2:1). The product was isolated in quantitative yield. Mp 129 - 130
- 35 °C.

1-(3-Nitrophenyl)-homopiperazine fumaric acid salt (14J)

A mixture of 1-fluoro-3-nitrobenzene (10.0 g, 71 mmol), and homopiperazine (21.3 g, 213 mmol) in dioxane (100 ml) was refluxed for 15 h. Water (100 ml) was added and the mixture
5 was extracted twice with ethyl acetate (100 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound. Yield 0.69 g, 4 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 163.1 - 164.4 °C.

10 1-(3-Aminophenyl)-homopiperazine fumaric acid salt (15J)

A mixture of 1-(3-nitrophenyl)-homopiperazine (2.4 g, 12.2 mmol), palladium on carbon (0.25 g, 5 %) and ethanol (75 ml) was stirred under hydrogen for 24 h. The crude mixture was filtered through celite. Aqueous sodium hydroxide (50 ml, 1 M) was added and the mixture was extracted twice with diethyl ether (50 ml). Yield 0.61 g, 26 %. The corresponding salt was
15 obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 165.4 - 167.7 °C.

1-(3-Methoxyphenyl)-homopiperazine fumaric acid salt (16J)

A mixture of 3-bromoanisole (10.0 g, 53.4 mmol), homopiperazine (10.7 g, 106.9 mmol),
20 potassium *tert*-butoxide (10.6 g, 106.9 mmol), tetrakis(triphenylphosphine)palladium(0) (62 mg, 0.53 mmol) and 1,2-dimethoxyethane (100 ml) was stirred at reflux for 1 h. tetrakis(triphenylphosphine)palladium(0) (286 mg, 0.247 mmol) Sodium hydroxide (1 M, 100 ml) was added and the mixture was extracted two times with ethyl acetate (150 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1)
25 gave the title compound as free base. Yield 0.79 g, 7 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 164.1 - 165.7 °C.

1-(3-Hydroxyphenyl)-homopiperazine fumaric acid salt (17J)

30 Boron tribromide (11.1g, 44.4 mmol) was added to a mixture of 1-(3-Methoxyphenyl)-homopiperazine (0.54 g, 2.8 mmol) in dichloromethane (50 ml) at -70 °C. The reaction was allowed to reach room temperature overnight. The precipitated crystals were filtered and purified by chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound as free base. Yield 0.22 g, 41 %. The corresponding salt

was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 165 °C, decomp.

General procedure for 3-bromo and 3-chloro-5- or 6-alkoxypyridines:

5

3-Bromo-5-methoxypyridine

Sodium (2.33 g, 101.3 mmol) was added to methanol (50 ml) and was allowed to react, the mixture was evaporated. 3,5-Dibromopyridine (20.0 g, 84.4 mmol) and dimethyl sulfoxide (100 ml) were added. The mixture was stirred at 90 °C for 2 h. Sodium hydroxide (400 ml, 1 M) was
10 added and the mixture was extracted twice with diethyl ether (200 ml). Chromatography on silica gel with dichloromethane and 3% ethanol as solvent gave the title compound. Yield 10.6 g, 67 %. Mp 30-32 °C.

General procedure for 3-bromo and 3-chloro-5-arylpyridines:

15

3-Bromo-5-phenylpyridine

A mixture of 3,5-dibromopyridine (10.0 g, 42.2 mmol), phenylboronic acid (4.6 g, 38.0 mmol), tetrakis(triphenylphosphine)palladium(0) (1.45 g, 1.25 mmol), potassium carbonate (17.5 g, 127 mmol), water (63 ml) and 1,2-dimethoxyethane (126 ml) was stirred at reflux overnight.
20 Aqueous sodium hydroxide (1 M, 60 ml) was added followed by extraction twice with diethyl ether (100 ml). Chromatography on silica gel with dichloromethane as solvent gave the title compound. Yield 6.1 g, 68 %, Mp 42 - 44 °C.

3-Bromo-6-thioethoxypyridine

25 A mixture of sodium thioethoxide (7.81 g, 92.9 mmol), 2,5-Dibromopyridine (20.0 g, 84.4 mmol) and dimethyl sulfoxide (100 ml). The mixture was stirred at 20 °C overnight. Sodium hydroxide (300 ml, 1 M) was added and the mixture was extracted twice with diethyl ether (200 ml). Chromatography on silica gel with dichloromethane : petroleum ether, (1:2) as eluent, gave the title compound as an oil. Yield 16.8 g, 85 %.

30

3-Bromo-5-thioethoxypyridine

Prepared according to 3-Bromo-6-thioethoxypyridine, using 40 °C as reaction temperature. The title compound was obtained as an oil

35 **Method K**

1-(6-Chloro-3-pyridazinyl)-homopiperazine fumaric acid salt (1K)

A mixture of 3,6-dichloropyridazine (5.0 g, 33.5 mmol), homopiperazine (3.36 g, 33.5 mmol) and 50 ml of toluene was stirred at reflux for 0.5 h. Sodium hydroxide (50 ml, 1 M) was added and the mixture was extracted three times with ethyl acetate (100 ml). Chromatography on
5 silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound as free base. Yield 2.2 g, 31 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 165 - 166 °C.

1-(6-Phenyl-3-pyridazinyl)-homopiperazine fumaric acid salt (2K)

10 Was prepared according to method K from 2-chloro-6-phenylpyridazine. Mp 187 - 189 °C.

1-(6-Chloro-2-pyrazinyl)-homopiperazine fumaric acid salt (3K)

Was prepared according to method K from 2,6-dichloropyrazine. Mp 180 - 181 °C.

1-(3,6-Dimethyl-2-pyrazinyl)-homopiperazine fumaric acid salt (4K)

Was prepared according to method K, in the absence of solvent from 2-chloro-3,6-dimethylpyrazine at 130 °C. Mp 149 - 151 °C.

1-(6-Methyl-3-pyridazinyl)-homopiperazine fumaric acid salt (5K)

20 Was prepared according to method K in the absence of solvent at 130 °C. from 3-chloro-6-methyl-pyridazine. Mp 102 - 105 °C.

1-(5-Trifluoromethyl-3-pyridyl)-homopiperazine fumaric acid salt (6K)

Was prepared according to method K in the absence of solvent at 140 °C overnight from 3-
25 chloro-5-trifluoromethylpyridine. Mp 164 - 166 °C.

1-(3-Chloro-2-quinoxaliny)-homopiperazine fumaric acid salt (7K)

Was prepared according to method K in the absence of solvent at 130 °C for 4 h from 2,3-dichloroquinoxaline. Mp 136.2 - 139.9 °C.

30

1-(6-Bromo-3-pyridyl)-homopiperazine fumaric acid salt (8K)

1-(3-Pyridyl)-homopiperazine (0.885 g, 5.0 mmol) was solved in acetonitrile (50 ml). N-Bromosuccinimide (1.7 g, 10.0 mmol) was added, and the mixture was stirred for 15 min. The crude mixture was evaporated. Chromatography on silica gel with dichloromethane, methanol
35 and conc. ammonia (89:10:1) gave the title compound as free base. Yield 0.50 g, 39 %. The

corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 164 -166 °C.

1-(6-Bromo-3-pyridyl)-4- *tert*-butoxycarbonyl homopiperazine

5 1-(3-Pyridyl)-4- *tert*-butoxycarbonyl homopiperazine (25.0 g, 90.1 mmol) was solved in acetonitrile (400 ml) and cooled to 0 °C. N-Bromosuccinimide (19.3 g, 108.2 mmol) was added during 10 min. Water (400 ml) was added and the mixture was extracted twice with diethyl ether (200 ml). Chromatography on silica gel with a mixture of petroleum: ethyl acetate (1 : 1) gave the title compound as free base. Yield 18.7 g, 58 %.

10

1-(6-Chloro-3-pyridyl)-4- *tert*-butoxycarbonyl homopiperazine

A stirred solution of 1-(6-Bromo-3-pyridyl)-4- *tert*-butoxycarbonyl homopiperazine (3.6 g, 10 mmol) in tetrahydrofuran (200 ml) was cooled to -78 °C. *Tert*-butyllithium (14.7 ml, 1.5 M) in pentane was added during 10 min, followed by stirring for 5 min at -78 °C. 1,3-Dichloro-5,5-
15 dimethylhydantoin (1.97 g, 10 mmol) was added in small portions. The mixture was stirred for 0.5 min. Aqueous sodium hydroxide was added (100 ml, 4 M), and the reaction was allowed reach room temperature. The mixture was extracted twice with diethyl ether (100 ml) Chromatography on silica gel with a mixture of petroleum: ethyl acetate (1 : 1) gave the title compound as free base. Yield 1.7 g, 55 %.

20

1-(6-Chloro-3-pyridyl)-homopiperazine (9K)

A mixture of 1-(6-chloro-3-pyridyl)-4- *tert*-butoxycarbonyl homopiperazine (1.7 g, 5.5 mmol), trifluoro acetic acid (4.5 ml, 55 mmol) and dichloromethane at room temperature for 1 h. The mixture was evaporated. Chromatography on silica gel with a mixture of petroleum: ethyl
25 acetate (1 : 1) gave the title compound as free base. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9 : 1) saturated with fumaric acid. Mp 165 - 167 °C. Yield 0.72 g, 40 %.

1-[6-(N-Pyrrolidiny)-3-pyridyl]-4- *tert*-butoxycarbonyl homopiperazine

30 A stirred solution of 1-(6-Bromo-3-pyridyl)-4- *tert*-butoxycarbonyl homopiperazine (0.71 g, 2.0 mmol), pyrrolidine (2.0 g, 20 mmol), potassium *tert*-butoxide (0.45 g, 4.0 mmol), tetrakis(triphenylphosphine)palladium(0) (0.12 g, 0.10 mmol) and 1,2-dimethoxyethane was stirred at reflux overnight. Water (40 ml) was added and the mixture was extracted twice with ethyl acetate (20 ml) Chromatography on silica gel with a mixture of petroleum: ethyl acetate
35 (1 : 1) gave the title compound as free base. Yield 0.21 g, 30 %.

1-(6-phenyl-3-pyridyl)-4- *tert*-butoxycarbonyl homopiperazine (10K)

1-(6-Bromo-3-pyridyl)-4- *tert*-butoxycarbonyl homopiperazine (0.89 g, 2.5 mmol), phenyl boronic acid (0.37 g, 3.0 mmol), tetrakis(triphenylphosphine)palladium(0) (0.15 g, 0.12 mmol),
5 1,2-dimethoxyethane (50 ml), potassium carbonate (1.0 g, 7.5 mmol) and water (7.5 ml) was stirred at reflux overnight. Water (50 ml) was added and the mixture was extracted twice with ethyl acetate (40 ml). Chromatography on silica gel with a mixture of petroleum: ethyl acetate (1 : 1) gave the title compound as an oil. Yield 0.83 g, 94 %.

10 1-(6-Bromo-5-ethoxy-3-pyridyl)-4- *tert*-butoxycarbonyl homopiperazine

N-Bromosuccinimide (2.7g, 15.2 mmol) was added to 1-(5-Ethoxy-3-pyridyl)-4- *tert*-butoxycarbonyl homopiperazine (4.5 g, 14.0 mmol) at room temperature. The mixture was stirred for 2 min. The mixture was washed with saturated sodium sulfite (100 ml) was. Chromatography on silica gel with a mixture of petroleum: ethyl acetate (1 : 1) gave the title
15 compound as an oil. Yield 3.3 g, 58 %.

Method L**1-(3-Pyridazinyl)-homopiperazine (1L)**

A mixture of 1-(3-Chloro-6-pyridazinyl)-homopiperazine (5.56 g, 26.1 mmol), palladium on
20 carbon (2.1 g, 10 %) and ethanol (150 ml) was stirred under hydrogen overnight. The crude product was filtered through celite and evaporated. Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound as free base. Yield 2.78 g, 60 %, 185.0 - 186.9 °C.

25 1-(2-Quinoxaliny)-homopiperazine fumaric acid salt (2L)

Was prepared according to method L from 1-(3-Chloro-2-quinoxaliny)-homopiperazine Mp 177- 180 °C.

N-(5-Methoxy-3-pyridyl)-ethylenediamine (3L)

30 Sodium (4.98 g, 216.7 mmol) was added to methanol (100 ml), and was allowed to react, the mixture was evaporated. 3,5-Dichloropyridine (25.0 g, 166.7 mmol) and dimethyl sulfoxide (250 ml) were added. The mixture was stirred at 70 °C overnight. Aqueous sodium hydroxide (500 ml, 1 M) was added, and the mixture was extracted twice with diethyl ether (500 ml). The crude mixture was stirred together with ethylenediamine (50.0 g, 833.5 mmol), potassium *tert*-
35 butoxide (37.4 g, 333.4 mmol) and 1,2-dimethoxyethane (500 ml) at reflux for 4 h. Aqueous

sodium hydroxide (1 l, 1 M) was added, and the mixture was extracted five times with ethyl acetate (500 ml). Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound as free base. Yield 13.0 g, 47 %.

5 Method M

1-(5-Methoxy-3-pyridyl)-3-methyl-imidazolidine fumaric acid salt (1M)


A mixture of N-(5-methoxy-3-pyridyl)-ethylenediamine (0.76 g, 4.5 mmol), formic acid (6.3 g, 136.4 mmol) and formaldehyde (4.1 g, 136.4 mmol) was stirred at reflux for 2 h at reflux. The crude mixture was evaporated. Sodium hydroxide (50 ml, 4 M) was added and the mixture was
10 extracted with ethyl acetate. Chromatography on silica gel with dichloromethane, methanol and conc. ammonia (89:10:1) gave the title compound as free base. Yield 9.7 g, 44 %. The corresponding salt was obtained by addition of a diethyl ether and methanol mixture (9:1) saturated with fumaric acid. Mp 139 - 142 °C.

15 1-(5-Methoxy-3-pyridyl)-3-methyl-1.3-diazacyclohexane fumaric acid salt (2M)

Was prepared according to method M from N-(5-methoxy-3-pyridyl)-1,3-propylene-diamine. Mp 149 - 151 °C.

Claims:

1. A compound represented by the general formula



any of its enantiomers or any mixture thereof, isotopes thereof or a pharmaceutically acceptable salt thereof;

wherein

n is 1, 2 or 3;

m is 0, 1 or 2;

R represents hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, or aralkyl; and

R¹ represents **aminophenyl; nitrophenyl; hydroxyphenyl, alkoxyphenyl;**

a monocyclic 5 to 6 membered heterocyclic group which may be substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino, nitro, -COOR³, -CONR²R³, -NH-CO₂R², NHCO-R², -OCO-NR²R³;

wherein R² and R³ independently represents hydrogen or alkyl;

aryl optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

-X-alkyl-Y-alkyl wherein X and Y independently represents O, S, NH, N-alkyl or Se; and alkyl is optionally substituted with alkoxy or thioalkoxy;

-X-(alkyl)_o-aryl wherein o is 0 or 1 and X represents O, S, NH, N-alkyl or Se; optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

-X-(alkyl)_o-Z wherein o is 0 or 1 and X represents O, S, NH, N-alkyl or Se and Z represents a 5- or 6- membered monocyclic heterocyclic group; optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

- 5 - a monocyclic 5 to 6 membered heterocyclic group optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

or R¹ represents a **bicyclic heterocyclic group**, composed of a 5 to 6 membered monocyclic
 10 heterocyclic group fused to a benzene ring, and which may be substituted one or more times with substituents selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, alkoxy-alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino, nitro, aryl optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy,
 15 halogen, CF₃, OCF₃, CN, amino and nitro; and a monocyclic 5 to 6 membered heterocyclic group optionally substituted one or more times with alkyl, cycloalkyl, cycloalkylalkyl alkenyl, alkynyl, alkoxy, cycloalkoxy, alkenoxy, alkynoxy, methylenedioxy, halogen, CF₃, OCF₃, CN, amino and nitro;

- 20 with the proviso that when R¹ is hydroxyphenyl, alkoxyphenyl, nitrophenyl or aminophenyl, then m and n are not both 2;

2. A compound of formula (I), according to claim 1, wherein
 R and R¹ is attached to a piperazine, a homopiperazine, a 1,4-di-azacyclooctane, a 1,5-di-
 25 azacyclooctane, a 1,3 diazacyclohexane or an imidazolidine;

3. A compound of formula (I), according to claim 1, wherein R¹ represents **isoquinolinyl**;
 or
phenyl optionally substituted with nitro, hydroxy, amino or alkoxy; or
 30 **pyridyl, quinolinyl, pyridazinyl or pyridinyl** all of which may be optionally substituted one or more times with alkoxy, cycloalkoxy, alkoxyalkoxy, alkoxycycloalkyl, hydroxyalkoxy, thioalkoxy, thioalkylaryl, alkenoxy, alkynoxy, carboxylamido, a bicyclic heterocyclic group, thioalkoxyaryl, thioaryl, hydroxy or trifluoromethanesulfonyloxy; halogen, phenyl optionally substituted with nitro; a monocyclic 5 to 6 membered heterocyclic group optionally substituted with alkyl;

4. A compound of formula (I), according to claim 1, wherein R¹ represents 3-pyridyl, 3-quinolinyl, 5-methoxy-3-pyridyl, 5-chloro-3-pyridyl, 5-(4-methyl-1-piperazinyl)-3-pyridyl, 5-ethoxy-3-pyridyl, 6-methoxy-3-pyridyl, 5-propyloxy-3-pyridyl, 5-phenyl-3-pyridyl, 5-(2-methyl-propyloxy)-3-pyridyl, 5-propyl-1-en-oxy-3-pyridyl, 2-chloro-5-methoxy-3-pyridyl, 6-chloro-5-methoxy-3-pyridyl, 6-bromo-5-ethoxy-3-pyridyl, 6-(N-pyrrolidinyl)-3-pyridyl, 6-phenyl-3-pyridyl, 5-(3-nitrophenyl)-3-pyridyl, 5-butoxy-3-pyridyl, 5-methoxyethoxy-3-pyridyl, 5-(2-methyl-propoxy)-3-pyridyl, 5-(2-hydroxy-ethoxy)-3-pyridyl, 5-(3-methyl-butoxy)-3-pyridyl, 5-cyclopropylmethoxy-3-pyridyl, 5-propyloxy-3-pyridyl, 5-hexyloxy-3-pyridyl, 5-cyclohexylmethoxy-3-pyridyl, 6-thioethoxy-3-pyridyl, 5-(2-ethoxy-ethoxy)-3-pyridyl, 5-pentyloxy-3-pyridyl, 5-heptyloxy-3-pyridyl, 5-(propyl-1-en-oxy)-3-pyridyl, 5-thiobenzyl-3-pyridyl, 5-carboxylamido-3-pyridyl, 5-thiophenyl-3-pyridyl, 5-methoxy-methoxy-3-pyridyl, 5-(3-pyridyl)-3-pyridyl, 5-(1-pyrrolyl)-3-pyridyl, 5-(1-indolyl)-3-pyridyl, 5,6-dimethoxy-3-pyridyl, 5-ethenyloxy-3-pyridyl, 5-cyclopentyloxy-3-pyridyl, 5-(ethoxy-d5)-3-pyridyl, 3-chloro-5-pyridyl, 3-bromo-5-pyridyl, 4-isoquinolinyl, 5-hydroxy-3-pyridyl, 5-trifluoromethanesulfonyl-oxy-3-pyridyl, 5-ethynyl-3-pyridyl, 3-nitrophenyl, 3-aminophenyl, 3-methoxyphenyl, 3-hydroxyphenyl, 6-chloro-3-pyridazinyl, 6-phenyl-3-pyridazinyl, 6-chloro-2-pyrazinyl, 3,6-dimethyl-2-pyrazinyl, 6-methyl-3-pyridazinyl, 5-trifluoromethyl-3-pyridyl, 6-bromo-3-pyridyl, 6-chloro-3-pyridyl or 3-pyridazinyl;

5. A compound of formula (I), according to claim 1, wherein R represents hydrogen, lower alkyl or aralkyl;

6. A compound of formula (I), according to claim 1, said compound being:

- 4-Methyl-1-(3-pyridyl)-piperazine;
- 4-Methyl-1-(3-pyridyl)-piperazine;
- 4-Methyl-1-(5-methoxy-3-pyridyl)-piperazine;
- 3,5-Bis-[4,4'-methyl-1,1'-piperazinyl]pyridine;
- 1-(5-Chloro-3-pyridyl)-4-Methyl-piperazine;
- 4-Methyl-1-(5-phenyl-3-pyridyl)-piperazine;
- 1-(5-Ethoxy-3-pyridyl)-4-methyl-piperazine;
- 1-(5-Butoxy-3-pyridyl)-4-methyl-piperazine;
- 4-Methyl-1-[5-(propyl-1-en-oxy)-3-pyridyl]-piperazine;
- 1-(5-Ethenyloxy-3-pyridyl)-4-methyl-piperazine;
- 1-(5-Methoxy-3-pyridyl)-4-methyl-1,5-diazacyclooctane;
- 1-(6-Chloro-3-pyridazinyl)-4-methyl-1,5-diazacyclooctane;

- 1-(3-Pyridyl)-piperazine;
1-(3-Pyridyl)-piperazine;
1-(5-Methoxy-3-pyridyl)-piperazine;
1-(5-Chloro-3-pyridyl)-piperazine;
5 1-(5-Phenyl-3-pyridyl)-piperazine;
1-(5-Methoxy-3-pyridyl)-1,5-diazacyclooctane;
1-(6-Chloro-3-pyridazinyl)-1,5-diazacyclooctane;
1-(6-Chloro-3-pyridazinyl)-1,4-diazacyclooctane;
1-(5-Ethoxy-3-pyridyl)piperazine;
10 1-(5-Butoxy-3-pyridyl)piperazine;
1-[5-(Propyl-1-en-oxy)-3-pyridyl]-piperazine;
1-(5-Ethenyloxy-3-pyridyl)piperazine;
1-(5-Ethoxy-3-pyridyl)-1,5-diazacyclooctane,
1-(5-Propyloxy-3-pyridinyl)-1,5-diazacyclooctane,
15 1-[5-(Propyl-1-en-oxy)-3-pyridyl]-1,5-diazacyclooctane,
1-(5-Ethenyloxy-3-pyridyl) 1,5-diazacyclooctane;
1-(5-Ethoxy-3-pyridyl)-4-ethyl-piperazine;
4-Methyl-1-(3-pyridyl)-homopiperazine;
4-Methyl-1-(3-pyridyl)-homopiperazine;
20 1-(5-Methoxy-3-pyridyl)-4-methyl-homopiperazine;
1-(5-Ethoxy-3-pyridyl)-4-methyl-homopiperazine;
4-Methyl-1-(5-phenyl-3-pyridyl)-homopiperazine;
1-(5-Butoxy-3-pyridyl)-4-methyl-homopiperazine;
1-(5-Methoxyethoxy-3-pyridyl)-4-methyl-homopiperazine;
25 4-Methyl-1-[5-(2-methyl-propyloxy)-3-pyridyl]-homopiperazine;
1-(5-Cyclopropylmethoxy-3-pyridyl)-4-methyl-homopiperazine;
4-Methyl-1-(5-propyloxy-3-pyridyl)-homopiperazine;
1-(5-Hexyloxy-3-pyridyl)-4-methyl-homopiperazine;
4-Methyl-1-[5-(3-methyl-butoxy)-3-pyridyl]-homopiperazine;
30 4-Methyl-1-(6-thioethoxy-3-pyridyl)-homopiperazine;
1-(5-Cyclohexylmethoxy-3-pyridyl)-4-methyl-homopiperazine;
4-Methyl-1-(5-pentyloxy-3-pyridyl)-homopiperazine;
1-(5-Heptyloxy-3-pyridyl)-4-methyl-homopiperazine;
4-Methyl-1-(5-propyl-1-en-oxy-3-pyridyl)-homopiperazine;
35 4-Methyl-1-(5-thiobenzyl-3-pyridyl)-homopiperazine;

- 4-Methyl-1-[5-(3-pyridyl)-3-pyridyl]-homopiperazine;
4-Methyl-1-(3-nitrophenyl)-homopiperazine;
4-Methyl-1-(6-chloro-3-pyridazinyl)-homopiperazine;
4-Methyl-1-(6-phenyl-3-pyridazinyl)-homopiperazine;
5 4-Methyl-1-(3-pyridazinyl)-homopiperazine;
4-Methyl-1-(6-methyl-3-pyridazinyl)-homopiperazine;
1-(5-Cyclopentyloxy-3-pyridyl)-4-methyl-homopiperazine;
4-Benzyl-1-(3-pyridyl)-homopiperazine;
4-Ethyl-1-(3-pyridyl)-homopiperazine;
10 1-(3-Pyridyl)-homopiperazine;
1-(3-Pyridyl)-homopiperazine;
1-(6-Methoxy-3-pyridyl)-homopiperazine;
1-(2-Chloro-5-methoxy-3-pyridyl)-homopiperazine;
1-(6-Chloro-5-methoxy-3-pyridyl)-homopiperazine;
15 1-(6-Bromo-5-ethoxy-3-pyridyl)-homopiperazine;
1-[6-(N-Pyrrolidinyl)-3-pyridyl]-homopiperazine;
1-(6-Phenyl-3-pyridyl)-4-homopiperazine;
1-[5-(3-Nitrophenyl)-3-pyridyl]-4-homopiperazine;
1-(5-Methoxy-3-pyridyl)-homopiperazine;
20 1-(5-Phenyl-3-pyridyl)-homopiperazine;
1-(5-Ethoxy-3-pyridyl)-homopiperazine;
1-(5-Butoxy-3-pyridyl)-homopiperazine;
1-(5-Methoxyethoxy-3-pyridyl)-homopiperazine;
1-[5-(2-Methyl-propoxy)-3-pyridyl]-homopiperazine;
25 1-[5-(2-Hydroxy-ethoxy)-3-pyridyl]-homopiperazine;
1-[5-(3-Methyl-butoxy)-3-pyridyl]-homopiperazine;
1-(5-Cyclopropylmethoxy-3-pyridyl)-homopiperazine;
1-(5-Propyloxy-3-pyridyl)-homopiperazine;
1-(5-Hexyloxy-3-pyridyl)-homopiperazine;
30 1-(5-Cyclohexylmethoxy-3-pyridyl)-homopiperazine;
1-(6-Thioethoxy-3-pyridyl)-homopiperazine;
1-[5-(2-Ethoxy-ethoxy)-3-pyridyl]-homopiperazine;
1-(5-Pentyloxy-3-pyridyl)-homopiperazine;
1-(5-Heptyloxy-3-pyridyl)-homopiperazine;
35 1-[5-(Propyl-1-en-oxy)-3-pyridyl]-homopiperazine;

- 1-(5-Thiobenzyl-3-pyridyl)-homopiperazine;
1-(5-Carboxylamido-3-pyridyl)-homopiperazine;
1-(5-Thiophenyl-3-pyridyl)-homopiperazine;
1-[(5-Methoxy-methoxy)-3-pyridyl]-homopiperazine;
5 1-[5-(3-Pyridyl)-3-pyridyl]-homopiperazine;
1-(5-(1-Pyrrolyl)-3-pyridyl)-homopiperazine;
1-(5-(1-Indolyl)-3-pyridyl)-homopiperazine;
1-(5,6-Dimethoxy-3-pyridyl)-homopiperazine;
1-(5-Ethenyloxy-3-pyridyl)homopiperazine;
10 1-(5-Cyclopentyloxy-3-pyridyl)homopiperazine;
1-[5-(Ethoxy-d5)-3-pyridyl]-homopiperazine;
1-(3-Chloro-5-pyridyl)homopiperazine;
1-(3-Bromo-5-pyridyl)homopiperazine;
1-(4-Isoquinoliny)-homopiperazine;
15 1-(5-Hydroxy-3-pyridyl)-homopiperazine;
1-(5-Trifluoromethanesulfonyl-oxy-3-pyridyl)-homopiperazine;
1-(5-Ethynyl-3-pyridyl)-homopiperazine;
1-(3-Nitrophenyl)-homopiperazine;
1-(3-Aminophenyl)-homopiperazine;
20 1-(3-Methoxyphenyl)-homopiperazine;
1-(3-Hydroxyphenyl)-homopiperazine;
1-(6-Chloro-3-pyridazinyl)-homopiperazine;
1-(6-Phenyl-3-pyridazinyl)-homopiperazine;
1-(6-Chloro-2-pyrazinyl)-homopiperazine;
25 1-(3,6-Dimethyl-2-pyrazinyl)-homopiperazine;
1-(6-Methyl-3-pyridazinyl)-homopiperazine;
1-(5-Trifluoromethyl-3-pyridyl)-homopiperazine;
1-(6-Bromo-3-pyridyl)-homopiperazine;
1-(6-Chloro-3-pyridyl)-homopiperazine;
30 1-(3-Pyridazinyl)-homopiperazine;
N-(5-Methoxy-3-pyridyl)-ethylenediamine;
1-(5-Methoxy-3-pyridyl)-imidazolidine;
1-(5-Methoxy-3-pyridyl)-1,3-diazacyclohexane;
or a pharmaceutically acceptable addition salt thereof;

7. A pharmaceutical composition, comprising a therapeutically effective amount of a compound of any of the claims 1 to 6, or a pharmaceutically acceptable addition salt thereof, together with at least one pharmaceutically acceptable carrier or diluent.
- 5 8. A compound of any of the claims 1-6 for use as a medicament for treatment of a disease of a living animal body, including a human, which disease is responsive to the activity of nicotinic Ach receptors modulators;
9. The use of a compound according to any of the claims 1 to 6 for the manufacture of a
10 medicament for the treatment of a disease of a living animal body, including a human, which disease is responsive to the activity of nicotinic ACh receptor modulators.
10. The use according to claim 9 wherein the disease to be treated is pain, a disease in the central or peripheral system, a disease caused by smooth muscle contraction,
15 neurodegeneration, inflammation, chemical substance abuse or withdrawal symptoms caused by the cessation of intake of the chemical substance, such as assistance in the cessation of smoking;
11. The use of a compound according to claim 9 wherein a disease in the central or
20 peripheral system is Alzheimer's disease, Parkinson's disease, memory dysfunction or attention deficit hyperactivity disorder.
12. A method of treating a disease of a living animal body, including a human, which disease is responsive to the activity of nicotinic ACh receptor modulators, comprising the step
25 of administering to such a living animal body, including a human, in need thereof a therapeutically effective amount of a compound according to any of the claims 1 to 6.
13. A method according to claim 12, wherein pain, a disease in the central or peripheral system, a disease caused by smooth muscle contraction, neurodegeneration, inflammation,
30 chemical substance abuse or withdrawal symptoms caused by the cessation of intake of chemical substances, such as smoking cessation, is treated by administering to a mammal in need thereof a therapeutically effective amount of a compound of any of the claims 1-6;
14. The method according to claim 12 wherein a disease in the central or peripheral
35 system, said disease being Alzheimer's disease, Parkinson's disease, memory dysfunction or

attention deficit hyperactivity disorder, is treated by administering to a mammal in need thereof a therapeutically effective amount a compound of any of the 1-6;

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 98/00465

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D213/74 A61K31/495 C07D401/04 C07D403/04 C07D243/08
C07D401/14 C07D405/04 C07D409/04 C07D409/14 C07D417/04
C07D417/14 C07D215/38

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL ABSTRACTS, vol. 89, no. 5, 31 July 1978 Columbus, Ohio, US; abstract no. 36473, GEORGIEV ET AL.: "Study of the effect of piperazine ..." XP002097079 & Farmatsiya 1977, 27(3),23-28 see abstract ---	1-14
X	US 4 179 563 A (BUTLER DONALD E) 18 December 1979 see column 4; example 21 ---	1,2,5, 7-14
X	EP 0 361 489 A (CHUGAI PHARMACEUTICAL CO LTD) 4 April 1990 see the whole document ---	1,2,5, 7-14
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

18 March 1999

Date of mailing of the international search report

12.04.99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Steendijk, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 98/00465

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 117, no. 17, 26 October 1992 Columbus, Ohio, US; abstract no. 171353, XU ET AL.: "Synthesis and anticonvulsant activity..." XP002097080 & Beijing Yike Daxue Xuebao, 1991, 23(9), 477-480 see abstract	1-5, 7-10, 12, 13
X	GB 1 492 528 A (MERCK & CO INC) 23 November 1977 see the whole document	1-5, 7-10, 12, 13
X	US 4 163 849 A (LUMMA WILLIAM C JR ET AL) 7 August 1979 see the whole document	1-5, 7-10, 12, 13
X	EP 0 156 433 A (JANSSEN PHARMACEUTICA NV) 2 October 1985 see the whole document	1-5, 7, 8
X	WO 96 10568 A (MERCK & CO INC ; CHIANG YUAN CHING P (US); FINKE PAUL E (US); MACCO) 11 April 1996 see claim 1	1, 2, 5, 7-10, 12, 13
X	EP 0 385 350 A (NISSHIN FLOUR MILLING CO) 5 September 1990 see the whole document	1, 2, 5, 7, 8
X	EP 0 211 457 A (JANSSEN PHARMACEUTICA NV) 25 February 1987 see examples and intermediate IV	1-6
X	EP 0 802 173 A (TOSOH CORP) 22 October 1997 see examples 1, 8	1-6
X	US 4 806 536 A (CROSS PETER E ET AL) 21 February 1989 see intermediate IV	1-6
X	WO 94 18978 A (JANSSEN PHARMACEUTICA NV ; HEERES JAN (BE); MOSTMANS JOSEPH HECTOR) 1 September 1994 see page 21, line 14	1-5
	-/--	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 98/00465

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 66, no. 9, 27 February 1967 Columbus, Ohio, US; abstract no. 37889, HENRY: "A facile synthesis of piperazines..." XP002097081 & J.Heterocycl.Chem. 3(4), 503-511 (1966) see abstract ---	1-5
X	CHEMICAL ABSTRACTS, vol. 67, no. 11, 11 September 1967 Columbus, Ohio, US; abstract no. 54102, POPOV: "Synthesis of 1,4-unsymmetrically substituted piperazines..." XP002097082 & C.R.Acad.Bulg.Sci. 19(12), 1163-1166 see abstract ---	1-5
X	CHEMICAL ABSTRACTS, vol. 95, no. 9, 31 August 1981 Columbus, Ohio, US; abstract no. 80681, SCHOENEN: "Synthesis and reactions of 3-aminoquinoline" XP002097083 & Arch.Pharm. 314(5), 464-470 (1981) see abstract ---	1-5
E	WO 98 56792 A (BACH ALFRED ;BASF AG (DE); DULLWEBER UTA (DE); WICKE KARSTEN (DE);) 17 December 1998 see page 9 -----	1-5

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 98/00465

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-3,5 (all partially)
because they relate to subject matter not required to be searched by this Authority, namely:
The number of known compounds falling under the scope of the partially searched claims precludes citation of all prior art. The search has been guided by exemplified structures and the alledged activity
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 98/00465

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4179563 A	18-12-1979	BE 876389 A CA 1109468 A DE 2920250 A FR 2426051 A GB 2021113 A,B JP 55000384 A	17-09-1979 22-09-1981 22-11-1979 14-12-1979 28-11-1979 05-01-1980
EP 0361489 A	04-04-1990	FI 894629 A JP 2167265 A US 5019574 A	31-03-1990 27-06-1990 28-05-1991
GB 1492528 A	23-11-1977	AR 210349 A AT 353795 B AU 496854 B BE 840904 A BG 34185 A CA 1059128 A CH 619462 A CS 195726 B CY 1090 A DD 124599 A DE 2617205 A DK 164476 A,B, EG 12387 A FI 760978 A,B, FR 2308367 A GR 59900 A HK 60380 A IE 42979 B JP 1030870 C JP 51136688 A JP 55022475 B KE 3088 A LU 74795 A NL 7603800 A,B, PT 65027 B SE 421695 B SE 7604093 A US 4081542 A ZA 7602320 A	29-07-1977 10-12-1979 02-11-1978 16-08-1976 15-07-1983 24-07-1979 30-09-1980 29-02-1980 27-12-1980 02-03-1977 28-10-1976 22-10-1976 31-12-1978 22-10-1976 19-11-1976 16-03-1978 07-11-1980 19-11-1980 29-01-1981 26-11-1976 17-06-1980 07-11-1980 07-02-1977 25-10-1976 03-04-1978 25-01-1982 22-10-1976 28-03-1978 30-11-1977
US 4163849 A	07-08-1979	NONE	
EP 0156433 A	02-10-1985	US 5001125 A AT 61050 T AU 576563 B AU 4034885 A BG 43690 A CA 1238321 A CS 8501952 A DK 134185 A,B, FI 851177 A,B GR 850714 A JP 60226862 A PT 80157 B SU 1384198 A US 5157035 A US 5292738 A	19-03-1991 15-03-1991 01-09-1988 03-10-1985 15-07-1988 21-06-1988 18-11-1992 27-09-1985 27-09-1985 14-06-1985 12-11-1985 20-03-1987 23-03-1988 20-10-1991 08-03-1994

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 98/00465

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9610568 A	11-04-1996	US 5607936 A AU 3642995 A CA 2199621 A EP 0783498 A JP 10508297 T	04-03-1997 26-04-1996 11-04-1996 16-07-1997 18-08-1998
EP 0385350 A	05-09-1990	DE 69014351 D DE 69014351 T KR 130758 B US 4994456 A CA 2011143 A JP 2843632 B JP 3007258 A	12-01-1995 20-04-1995 09-04-1998 19-02-1991 01-09-1990 06-01-1999 14-01-1991
EP 0211457 A	25-02-1987	AU 6073986 A DK 363186 A FI 863120 A GR 862022 A JP 62029575 A PT 83098 B	05-02-1987 01-02-1987 01-02-1987 24-12-1986 07-02-1987 13-07-1988
EP 0802173 A	22-10-1997	JP 10139742 A JP 10081667 A	26-05-1998 31-03-1998
US 4806536 A	21-02-1989	AT 94534 T AU 575740 B AU 6858887 A CA 1303042 A DE 3787384 D DE 3787384 T DK 60887 A,B, EP 0233051 A ES 2058105 T FI 870499 A,B IE 60300 B JP 6070015 B JP 62185073 A PT 84248 B	15-10-1993 04-08-1988 13-08-1987 09-06-1992 21-10-1993 13-01-1994 05-10-1987 19-08-1987 01-11-1994 08-08-1987 29-06-1994 07-09-1994 13-08-1987 14-09-1989
WO 9418978 A	01-09-1994	AP 435 A AU 675508 B AU 6108194 A CA 2155830 A CN 1118138 A EP 0684821 A FI 953899 A HU 73662 A IL 108703 A JP 8506821 T LT 1874 A,B MX 9401294 A NO 953254 A NZ 261854 A PL 310348 A US 5650411 A ZA 9401137 A	17-11-1995 06-02-1997 14-09-1994 01-09-1994 06-03-1996 06-12-1995 18-08-1995 30-09-1996 04-01-1998 23-07-1996 25-09-1994 31-08-1994 11-09-1995 24-04-1997 11-12-1995 22-07-1997 18-08-1995
WO 9856792 A	17-12-1998	DE 19724980 A	17-12-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 98/00465

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9856792 A		AU 7917698 A	30-12-1998